

Partial wave decomposition of pion and photoproduction amplitudes

A. Anisovich^{1,2}, E. Klempt¹, A. Sarantsev^{1,2}, and U. Thoma³

¹ HISKP, Universität Bonn, D-53115 Bonn,

² Petersburg Nuclear Physics Institute, Gatchina, Russia

³ Physikalisches Institut, Universität Giessen, Germany

July 20, 2004

Abstract. Partial wave amplitudes for production and decay of baryon resonances are constructed in the framework of the operator expansion method. The approach is fully relativistically invariant and allow us to perform combined analyses of different reactions imposing directly analyticity and unitarity constraints. All formulas are given explicitly in the form used by the Crystal Barrel collaboration in the (partly forthcoming) analyses of the electro-, photo- and pion induced meson production data.

1 Introduction

The perturbative approach to the theory of strong interaction (perturbative QCD) cannot be applied directly to the region of low and intermediate energies. In spite of many efforts to create a non-perturbative formulation for QCD from first principles, a final breakthrough has not yet been achieved even if recent results of lattice QCD indicate that this situation might change in the future. A necessary step towards a better understanding of strong interactions is undoubtedly a precise knowledge of the experimental situation and a correct classification of strongly interacting particles.

In meson spectroscopy, considerable progress had been made during the last ten years. A variety of experiments lead to the discovery of a large number of new meson states. In particular scalar states, very purely known 15 years ago, are now one of the most studied systems. As a result, it is now possible to investigate systematically the question if additional states expected from QCD like glueballs or hybrids hide in the observed meson spectrum. Although there is still no agreement on the classification of scalar states, the number of reliable classifications is reduced to quite small number (see [1–4] and references therein). We expect that the new GSI facility will help to resolve the remaining ambiguities completely.

A very important observation is that those meson resonances which can be interpreted as dominant $q\bar{q}$ states are lying on linear trajectories, not only against the total spin but also against their radial quantum number [5]. Excitingly, this seems to be true also for baryons [6]. Almost all known baryons lye on linear trajectories with the same slope as that for mesons.

Most information about baryons comes from pion- and photon-induced production of single mesons. However the experience from meson spectroscopy shows that excited states decay dominantly into multi-body channels and are not observed reliably in the elastic cross section. Thus reactions with three or more final states provide rich information about the properties of hadronic resonances. One of recent examples is the possible observation of a pentaquark [4] which up to now was seen only in reactions with three or more final-state particles.

The task to extract pole positions and residues from multi-body final states is however not a simple one. Main problems can be traced to the large interference effects between different isobars and to contributions from singularities related to multi-body interactions. In [7] an approach based on the dispersion N/D method was put forward and successfully applied to the analysis of meson resonances. In this method allowed singularities in the reaction can be classified, resonances which are closest to the physical region can be taken into account accurately, and other contributions can be parameterized in an efficient way.

One of the key points in this approach is the operator decomposition method which provides a tool for a universal construction of partial wave amplitudes for reactions with two- and many-body final states. The operator decomposition method has a long history. It was used for the analysis of the reactions with three particle final states already in [8]. The full description of the method for the nonrelativistic case was given in [9]. The full relativistic approach for $NN \rightarrow NN(N\Delta)$ and $\gamma d \rightarrow pn$ was developed in [10–12]. The construction of partial wave amplitudes for production of meson resonances in different reactions can be found in [13–15].

In the present article we develop the operator expansion method to describe baryon resonances in meson- and

photon-induced reactions. The photon can be real or virtual, we assume it to be virtual unless the opposite is explicitly stated. The method is also very convenient to calculate contributions from triangle and box diagrams and to project t and u-channel exchange amplitudes into partial waves. The latter feature is very important for amplitudes near their unitarity limits where the unitarity property must be taken into account explicitly.

The formulas given here reproduce exactly the amplitudes used by the Crystal-Barrel-ELSA collaboration in their (partly forthcoming) analyses of single- and two-body photoproduction reactions.

It must be emphasized that a wealth of data on baryon resonances has been taken, is being analyzed or is going to be produced in the near future. At MAMI in Mainz [16] precision data were taken in the low-energy range which will be extended to 1.4 GeV photon energies in the close future. The GRAAL [17] experiment has produced invaluable data in particular using linearly polarized photons; the SAPHIR [18] experiment at Bonn has published a series of papers covering many basic photoproduction cross sections. The experiment is now replaced by the Crystal Barrel detector [19] which had produced before many results at the Low-Energy-Antiproton-Ring (LEAR) at CERN. And, last not least, Jlab at Newport News/Virginia [20] has accumulated high statistic data sets on photo- and electro-production of a variety of final states. First high-quality data have been published.

1.1 Orbital angular momentum operators

$$X_{\mu_1 \mu_2 \dots \mu_{L-1} \mu_L}^{(L)}$$

Let us consider a decay of a composite particle with spin J and momentum P ($P^2 = s$) into two spinless particles with momenta k_1 and k_2 . The only measured quantities in such a reaction are the particle momenta. The angular dependent part of the wave function of the composite state is described by operators constructed out of these momenta and the metric tensor. Such operators (we will denote them as $X_{\mu_1 \dots \mu_L}^{(L)}$) are called orbital angular momentum operators and correspond to irreducible representations of the Lorentz group. They satisfy the following properties [14]:

- Symmetry with respect to permutation of any two indices:

$$X_{\mu_1 \dots \mu_i \dots \mu_j \dots \mu_L}^{(L)} = X_{\mu_1 \dots \mu_j \dots \mu_i \dots \mu_L}^{(L)}. \quad (1)$$

- Orthogonality to the total momentum of the system, $P = k_1 + k_2$.

$$P_{\mu_i} X_{\mu_1 \dots \mu_i \dots \mu_L}^{(L)} = 0 \quad (2)$$

- The traceless property for summation over two any indices:

$$g_{\mu_i \mu_j} X_{\mu_1 \dots \mu_i \dots \mu_j \dots \mu_L}^{(L)} = 0. \quad (3)$$

Let us consider a one-loop diagram describing the decay of a composite system into two spinless particles which propagate and then form again a composite system. The decay and formation processes are described by orbital angular momentum operators. Due to conservation of quantum numbers this amplitude must vanish for initial and final states with different spin. The S-wave operator is a scalar and can be taken as unit operator. The P-wave operator is a vector. In the dispersion relation approach it is sufficient that the imaginary part of the loop diagram with S and P-wave operators as vertices is equal to 0. In the case of spinless particles this requirement entails

$$\int \frac{d\Omega}{4\pi} X_{\mu}^{(1)} = 0 \quad (4)$$

where the integral is taken over the solid angle of the relative momentum. In general the result of such an integration is proportional to the total momentum of the system P_{μ} (the only external vector):

$$\int \frac{d\Omega}{4\pi} X_{\mu}^{(1)} = \lambda P_{\mu} \quad (5)$$

Convoluting this expression with P_{μ} and demanding $\lambda = 0$ we obtain the orthogonality condition (2). The orthogonality between D-wave and S-wave is provided by the traceless condition (3); conditions (2,3) provide the orthogonality for all operators with different orbital angular momenta.

The orthogonality condition (2) is automatically fulfilled if the operators are constructed from the relative momenta k_{μ}^{\perp} and the tensor $g_{\mu\nu}^{\perp}$. They both are orthogonal to the total momentum of the system:

$$k_{\mu}^{\perp} = \frac{1}{2} g_{\mu\nu}^{\perp} (k_1 - k_2)_{\nu} \quad g_{\mu\nu}^{\perp} = g_{\mu\nu} - \frac{P_{\mu} P_{\nu}}{s} \quad (6)$$

In the center-of-mass system (c.m.s. from now onwards), where $P = (P_0, \mathbf{P}) = (\sqrt{s}, 0)$, the vector k^{\perp} is space-like: $k^{\perp} = (0, \mathbf{k})$.

The operator for spin $J = 0$ is a scalar (for example a unit operator), and the operator for spin $J = 1$ is a vector which can only be constructed from k_{μ}^{\perp} . The orbital angular momentum operators for $L = 1$ to 3 are:

$$\begin{aligned} X^{(0)} &= 1, & X_{\mu}^{(1)} &= k_{\mu}^{\perp}, \\ X_{\mu_1 \mu_2}^{(2)} &= \frac{3}{2} \left(k_{\mu_1}^{\perp} k_{\mu_2}^{\perp} - \frac{1}{3} k_{\perp}^2 g_{\mu_1 \mu_2}^{\perp} \right), \\ X_{\mu_1 \mu_2 \mu_3}^{(3)} &= \\ &= \frac{5}{2} \left[k_{\mu_1}^{\perp} k_{\mu_2}^{\perp} k_{\mu_3}^{\perp} - \frac{k_{\perp}^2}{5} (g_{\mu_1 \mu_2}^{\perp} k_{\mu_3}^{\perp} + g_{\mu_1 \mu_3}^{\perp} k_{\mu_2}^{\perp} + g_{\mu_2 \mu_3}^{\perp} k_{\mu_1}^{\perp}) \right]. \end{aligned} \quad (7)$$

The operators $X_{\mu_1 \dots \mu_L}^{(L)}$ for $L \geq 1$ can be written in form of a recurrent expression:

$$\begin{aligned} X_{\mu_1 \dots \mu_L}^{(L)} &= k_{\alpha}^{\perp} Z_{\mu_1 \dots \mu_L}^{\alpha}, \\ Z_{\mu_1 \dots \mu_L}^{\alpha} &= \frac{2L-1}{L^2} \left(\sum_{i=1}^L X_{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_L}^{(L-1)} g_{\mu_i \alpha}^{\perp} - \right. \end{aligned}$$

$$\frac{2}{2L-1} \sum_{\substack{i,j=1 \\ i < j}}^L g_{\mu_i \mu_j}^\perp X_{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_{j-1} \mu_{j+1} \dots \mu_L}^{(L-1)} \Big). \quad (8)$$

Convolution equality reads:

$$X_{\mu_1 \dots \mu_L}^{(L)} k_{\mu_L}^\perp = k_\perp^2 X_{\mu_1 \dots \mu_{L-1}}^{(L-1)}. \quad (9)$$

Based on eq.(9) and taking into account the traceless property of $X_{\mu_1 \dots \mu_L}^{(L)}$, one can write down the orthogonality-normalization condition for orbital angular operators:

$$\int \frac{d\Omega}{4\pi} X_{\mu_1 \dots \mu_n}^{(n)}(k^\perp) X_{\mu_1 \dots \mu_m}^{(m)}(k^\perp) = \delta_{nm} \alpha(L) k_\perp^{2n},$$

$$\alpha(L) = \prod_{l=1}^L \frac{2l-1}{l} = \frac{(2L-1)!!}{L!}. \quad (10)$$

Iterating eq. (8) one obtains the following expression for the operator $X_{\mu_1 \dots \mu_L}^{(L)}$:

$$X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) = \alpha(L) \left[k_{\mu_1}^\perp k_{\mu_2}^\perp k_{\mu_3}^\perp k_{\mu_4}^\perp \dots k_{\mu_L}^\perp - \frac{k_\perp^2}{2L-1} \left(g_{\mu_1 \mu_2}^\perp k_{\mu_3}^\perp k_{\mu_4}^\perp \dots k_{\mu_L}^\perp + g_{\mu_1 \mu_3}^\perp k_{\mu_2}^\perp k_{\mu_4}^\perp \dots k_{\mu_L}^\perp + \dots \right) + \frac{k_\perp^4}{(2L-1)(2L-3)} \left(g_{\mu_1 \mu_2}^\perp g_{\mu_3 \mu_4}^\perp k_{\mu_5}^\perp k_{\mu_6}^\perp \dots k_{\mu_L}^\perp + g_{\mu_1 \mu_2}^\perp g_{\mu_3 \mu_5}^\perp k_{\mu_4}^\perp k_{\mu_6}^\perp \dots k_{\mu_L}^\perp + \dots \right) + \dots \right]. \quad (11)$$

The angular part of the amplitude for scattering of two spinless particles (for example a $\pi\pi \rightarrow \pi\pi$ scattering) is described as a convolution of the operators $X^{(L)}(k)$ and $X^{(L)}(q)$ where k and q are relative momenta before and after the interaction.

$$X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) X_{\mu_1 \dots \mu_L}^{(L)}(q^\perp) = \alpha(L) \left(\sqrt{k_\perp^2} \sqrt{q_\perp^2} \right)^L P_L(z) \quad (12)$$

Here $P_L(z)$ are Legendre polynomials (see Appendix A) and $z = (k^\perp q^\perp) / (\sqrt{k_\perp^2} \sqrt{q_\perp^2})$ which are, in c.m.s., functions of the cosine of the angle between initial and final particles.

A comment: one should be careful with expression $\sqrt{k_\perp^2}$. In c.m.s.,

$$\sqrt{k_\perp^2} = \sqrt{-\mathbf{k}^2} = i|\mathbf{k}|$$

$$\left(\sqrt{k_\perp^2} \sqrt{q_\perp^2} \right)^L = (-1)^L (|\mathbf{k}||\mathbf{q}|)^L \quad (13)$$

1.2 The boson projection operator

Let us consider the imaginary part of the one-loop diagram when particles interact with relative momentum p , then

propagate with momentum k , and interact for a second time getting the relative momentum q . The process can be described by orbital angular momentum operators in the form

$$X_{\mu_1 \dots \mu_L}^{(L)}(p^\perp) \int \frac{d\Omega}{4\pi} X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) X_{\nu_1 \dots \nu_L}^{(L)}(k^\perp) X_{\nu_1 \dots \nu_L}^{(L)}(q^\perp)$$

The projection operator $O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}$ for the partial wave with angular momentum L is defined as:

$$\int \frac{d\Omega}{4\pi} X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) X_{\nu_1 \dots \nu_L}^{(L)}(k^\perp) = \frac{\alpha(L)}{2L+1} k_\perp^{2L} O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} \quad (14)$$

and satisfies the following relations:

$$X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp) O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} = X_{\nu_1 \dots \nu_L}^{(L)}(k^\perp),$$

$$O_{\alpha_1 \dots \alpha_L}^{\mu_1 \dots \mu_L} O_{\nu_1 \dots \nu_L}^{\alpha_1 \dots \alpha_L} = O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}. \quad (15)$$

Due to properties (15), the product of any number of loop diagrams will be described by the same projection operator. This operator has the same symmetry, orthogonality and traceless properties as X -operators (for the same set of up and down indices) but the O -operator does not depend on the relative momentum of the constituents and does not describe decay processes. It represents the propagation of the composite system and defines the structure of the boson propagator (its numerator). More details on the properties of X and O -operators can be found in [14].

Taking into account the definition of the projection operators (15) and the properties of the X -operators (12) we obtain:

$$k_{\mu_1} \dots k_{\mu_L} O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} = \frac{1}{\alpha(L)} X_{\nu_1 \dots \nu_L}^{(L)}(k^\perp). \quad (16)$$

This equation presents the basic property of the projection operator: it projects any operator with L indices onto the partial wave operator with angular momentum L . The projection operator can also be calculated using the recurrent expression:

$$O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} = \frac{1}{L^2} \left(\sum_{i,j=1}^L g_{\mu_i \nu_j}^\perp O_{\nu_1 \dots \nu_{j-1} \nu_{j+1} \dots \nu_L}^{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_L} - \frac{4}{(2L-1)(2L-3)} \sum_{\substack{i < j \\ k < m}}^L g_{\mu_i \mu_j}^\perp g_{\nu_k \nu_m}^\perp O_{\nu_1 \dots \nu_{k-1} \nu_{k+1} \dots \nu_{m-1} \nu_{m+1} \dots \nu_L}^{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_{j-1} \mu_{j+1} \dots \mu_L} \right) \quad (17)$$

The low order projection operators are:

$$O = 1 \quad O_\nu^\mu = g_{\mu\nu}^\perp$$

$$O_{\alpha\beta}^{\mu\nu} = \frac{1}{2} \left(g_{\mu\alpha}^\perp g_{\nu\beta}^\perp + g_{\mu\beta}^\perp g_{\nu\alpha}^\perp - \frac{2}{3} g_{\mu\nu}^\perp g_{\alpha\beta}^\perp \right) \quad (18)$$

1.3 The vector projection operator in the gauge invariant limit

The sum over the possible polarisations of a vector particle ε_μ with non zero mass corresponds to the vector projection operator:

$$\sum_\alpha \varepsilon_\mu^\alpha \varepsilon_\nu^{*\alpha} = O_\nu^\mu = g_{\mu\nu}^\perp \tag{19}$$

which means that there are three independent polarisation vectors orthogonal to the momentum of the particle and normalized as $\varepsilon_\mu^\alpha \varepsilon_\mu^{*\alpha} = -1$.

However photon polarisation vectors have only two independent components, its momentum squared is equal to 0 and therefore, the projection operator can not have the form (19). The invariant expression for the photon projection operator can be only constructed for the interaction of the photon with another particle. In this case it has the form:

$$g_{\mu\nu}^{\perp\perp} = -\sum_\alpha \varepsilon_\mu^\alpha \varepsilon_\nu^\alpha = g_{\mu\nu} - \frac{P_\mu P_\nu}{P^2} - \frac{k_\mu^\perp k_\nu^\perp}{k_\perp^2} \tag{20}$$

where k_1 is momentum of the baryon, k_2 -is momentum of the photon, $P = k_1 + k_2$ and

$$k_\mu^\perp = \frac{1}{2}(k_1 - k_2)_\nu g_{\mu\nu}^\perp = \frac{1}{2}(k_1 - k_2)_\nu \left(g_{\mu\nu}^\perp - \frac{P_\mu P_\nu}{P^2} \right) \tag{21}$$

In the c.m.s. with the momentum of γ being parallel to the z-axis, the $g_{\mu\nu}^{\perp\perp}$ tensor has a very simple form:

$$g_{\mu\nu}^{\perp\perp} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \tag{22}$$

where the vector components are defined as

$$p = (E, p_x, p_y, p_z).$$

The tensor (20) is orthogonal to the momentum of the both particles:

$$g_{\mu\nu}^{\perp\perp} k_{2\mu} = g_{\mu\nu}^{\perp\perp} k_{1\mu} = 0. \tag{23}$$

and it extracts the gauge invariant part of the amplitude. For the real photon:

$$A_\mu \varepsilon_\mu^\alpha = A_\nu g_{\nu\mu}^{\perp\perp} \varepsilon_\mu^\alpha \tag{24}$$

and the expression $A_\nu g_{\nu\mu}^{\perp\perp}$ is gauge invariant:

$$A_\nu g_{\nu\mu}^{\perp\perp} k_{2\mu} = 0.$$

2 Fermions

The wave function of a fermion is described as Dirac bispinor, as object in Dirac space represented by γ matrices. In the standard representation the γ matrices have the following form:

$$\gamma_0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma = \begin{pmatrix} 0 & \sigma \\ -\sigma & 0 \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \tag{25}$$

where σ are 2×2 Pauli matrices. In this representation the spinors for fermion particles with momentum p are:

$$u(p) = \frac{1}{\sqrt{p_0 + m}} \begin{pmatrix} (p_0 + m)\omega \\ (\mathbf{p}\sigma)\omega \end{pmatrix}, \quad \bar{u}(p) = \frac{((p_0 + m)\omega^*, -(\mathbf{p}\sigma)\omega^*)}{\sqrt{p_0 + m}}. \tag{26}$$

Here ω represents a 2-dimensional spinor and ω^* the conjugated and transposed spinor. The normalization condition can be written as:

$$\bar{u}(p)u(p) = 2m \sum_{polarizations} u(p)\bar{u}(p) = m + \hat{p} \tag{27}$$

We define $\hat{p} = p^\mu \gamma_\mu$.

3 The structure of fermion propagator

The wave function of a particle with spin $J = L + 1/2$ and momentum p is described by a tensor bispinor $\Psi_{\mu_1 \dots \mu_L}$: it is a tensor in Dirac space. As the tensor it satisfies the same properties as a boson wave function:

$$\begin{aligned} p_{\mu_i} \Psi_{\mu_1 \dots \mu_L} &= 0 \\ \Psi_{\mu_1 \dots \mu_i \dots \mu_j \dots \mu_L} &= \Psi_{\mu_1 \dots \mu_j \dots \mu_i \dots \mu_L} \\ g_{\mu_i \mu_j} \Psi_{\mu_1 \dots \mu_L} &= 0 \end{aligned} \tag{28}$$

In addition the fermion wave function must satisfy the following properties:

$$\begin{aligned} (\hat{p} - m) \Psi_{\mu_1 \dots \mu_L} &= 0 \\ \gamma_{\mu_i} \Psi_{\mu_1 \dots \mu_L} &= 0 \end{aligned} \tag{29}$$

Conditions (28), (29) define the structure of the fermion propagator (projection operator) which can be written in the following form:

$$F_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(p) = (m + \hat{p}) R_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} \tag{30}$$

Here $(m + \hat{p})$ corresponds to the propagator for a fermion with $J = 1/2$.

The operator $R_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}$ describes the tensor structure of the propagator. It is equal to 1 for $J = 1/2$ particle and is proportional to $g_{\mu\nu}^\perp - \gamma_\mu^\perp \gamma_\nu^\perp / 3$ for a particle with spin $J = 3/2$ ($\gamma_\mu^\perp = g_{\mu\nu}^\perp \gamma_\nu$).

The conditions (28) are identical for fermion and boson projection operators and therefore the fermion projection operator can be written as:

$$R_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} = O_{\alpha_1 \dots \alpha_L}^{\mu_1 \dots \mu_L} T_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L} O_{\nu_1 \dots \nu_L}^{\beta_1 \dots \beta_L} \tag{31}$$

The $T_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}$ operator can be expressed in a rather simple form since all symmetry and orthogonality conditions are imposed by O -operators. First, the T-operators are constructed only out of metrical tensors and γ -matrices. Second, a construction like $\gamma_{\alpha_i} \gamma_{\alpha_j}$

$$\gamma_{\alpha_i} \gamma_{\alpha_j} = \frac{1}{2} g_{\alpha_i \alpha_j} + \sigma_{\alpha_i \alpha_j}, \tag{32}$$

where $\sigma_{\alpha_i \alpha_j} = \frac{1}{2} (\gamma_{\alpha_i} \gamma_{\alpha_j} - \gamma_{\alpha_j} \gamma_{\alpha_i})$

gives zero if multiplied with an O -operator (The first term due to the traceless conditions and the second one due to symmetry properties). The only structures which can then be constructed are $g_{\alpha_i\beta_j}$ and $\sigma_{\alpha_i\beta_j}$. Moreover, taking into account the symmetry properties of the O -operators, the latter can be used as $\sigma_{\alpha_1\beta_1}$:

$$T_{\beta_1\dots\beta_L}^{\alpha_1\dots\alpha_L} = \frac{L+1}{2L+1} \left(g_{\alpha_1\beta_1} - \frac{L}{L+1} \sigma_{\alpha_1\beta_1} \right) \prod_{i=2}^L g_{\alpha_i\beta_i} \quad (33)$$

Here the coefficients are calculated to satisfy the conditions (29) for the fermion projection operator:

$$\gamma_{\mu_i} F_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L} = F_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L} \gamma_{\nu_j} = 0 \quad (34)$$

$$F_{\alpha_1\dots\alpha_L}^{\mu_1\dots\mu_L} F_{\nu_1\dots\nu_L}^{\alpha_1\dots\alpha_L} = F_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L} \quad (35)$$

It is not necessary to construct the T operator out of the metric tensors and σ -matrices orthogonal to the momentum of the particle. Orthogonality is imposed by O -operators. However, to use the same ingredients for all operators, it is easier to introduce this property directly, rewriting the T -operators as:

$$T_{\beta_1\dots\beta_L}^{\alpha_1\dots\alpha_L} = \frac{L+1}{2L+1} \left(g_{\alpha_1\beta_1}^\perp - \frac{L}{L+1} \sigma_{\alpha_1\beta_1}^\perp \right) \prod_{i=2}^L g_{\alpha_i\beta_i}^\perp, \quad (36)$$

$$\sigma_{\mu\nu}^\perp = \frac{1}{2} (\gamma_\mu^\perp \gamma_\nu^\perp - \gamma_\nu^\perp \gamma_\mu^\perp)$$

3.1 Fermion propagator for an unstable particle

The numerator of a stable particle propagator has a very simple structure in its c.m.s.:

$$m + \hat{P} = 2m \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (37)$$

Assume a resonance with an invariant mass \sqrt{s} ($P^2 = s$). To maintain the orthogonality condition for the operators one should replace $m \rightarrow \sqrt{s}$ in eq. (30). Then, for a resonance in its c.m.s.:

$$\sqrt{s} + \hat{P} = 2\sqrt{s} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (38)$$

Such a structure is divergent at large energies and it is reasonable to regularize it with the factor $2M/(2\sqrt{s})$ or simply with $1/(2\sqrt{s})$ to provide a correct asymptotical behavior. Therefore we use the following expression for the numerator of a resonance propagator:

$$F_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L}(P) = \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} R_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L} \quad (39)$$

4 πN scattering

Let us now construct vertices for the decay of a composite baryon system with momentum P into the πN final state

with relative momentum $k = 1/2(k_1 - k_2)$ (here k_1 is the nucleon momentum). A particle with spin $J^P = 1/2^-$ decays into the πN channel in an S-wave, hence the orbital angular momentum operator is a scalar, e.g. a unit operator. For the vertex we get:

$$\bar{u}(P)u(k_1). \quad (40)$$

Here $u(P)$ is a bispinor of the composite particle and $u(k_1)$ is the bispinor of the nucleon. A resonance with spin $3/2^+$ decays into πN with an orbital angular momentum $L = 1$ and the vertex must be a vector, constructed out of k_μ^\perp and γ_μ^\perp . However it is sufficient to take only k_μ^\perp : first, due to the properties (29) and second, due to the fact that the projection operator (numerator of the fermion propagator) will automatically provide the correct structure. Thus we obtain for the decay of particles with $J = (L + 1/2)$, $P = (-1)^{L+1} (1/2^-, 3/2^+, 5/2^-, 7/2^+, \dots)$ the expression

$$\bar{\Psi}_{\mu_1\dots\mu_L} X_{\mu_1\dots\mu_L}^{(L)}(k^\perp)u(k_1). \quad (41)$$

Let us call below this set of states where the total angular momentum is given by the orbital angular momentum plus $1/2$ as 'plus' or '+' states. 'Minus' or '-' states are defined analogously ($J = (L - 1/2)$, $P = (-1)^{L+1}$).

It is convenient to introduce vertex functions $N_{\mu_1\dots\mu_L}^\pm$ describing the decay of a resonance into a pseudoscalar meson and a nucleon. Then for '+' states:

$$\begin{aligned} & \bar{\Psi}_{\mu_1\dots\mu_L} N_{\mu_1\dots\mu_L}^+(k^\perp)u(k_1) \\ N_{\mu_1\dots\mu_L}^+(k^\perp) &= X_{\mu_1\dots\mu_L}^{(L)}(k^\perp) \end{aligned} \quad (42)$$

The angular dependent part of the $\pi N \rightarrow$ resonance $\rightarrow \pi N$ transition amplitude can be constructed in a very simple way: the vertex function describing the interaction of the meson and the nucleon convolutes with the intermediate state propagator and the decay vertex function:

$$\bar{u}_f \tilde{N}_{\mu_1\dots\mu_L}^\pm F_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L}(P) N_{\nu_1\dots\nu_L}^\pm u_i \quad (43)$$

Here \tilde{N}^\pm is the left-hand vertex function (with two particles joining to one resonance) which is different from the decay vertex function N^\pm by the ordering of γ -matrices. (This is important for $N_{\mu_1\dots\mu_L}^-$ vertices which will be given on the next page). If q and k are the relative momenta before and after interaction and k_1 and q_1 are the corresponding nucleon momenta, the amplitude for πN scattering via '+' states can be written as:

$$A = \bar{u}(k_1) X_{\mu_1\dots\mu_L}(k^\perp) F_{\nu_1\dots\nu_L}^{\mu_1\dots\mu_L}(P) X_{\nu_1\dots\nu_L}(q^\perp) u(q_1) BW_L^+(s) \quad (44)$$

where $BW_L^+(s)$ describes the energy dependence of the intermediate state propagator. It is given, e.g., by a Breit-Wigner amplitude, a K-matrix or an N/D expression.

Using equations (15) and (37) we obtain:

$$A = \bar{u}(k_1) \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} u(q_1) (\sqrt{k_\perp^2} \sqrt{q_\perp^2})^L \frac{L+1}{2L+1} \alpha(L)$$

$$P_L(z)BW_L^+(s) - \bar{u}(k_1)\frac{\sqrt{s} + \hat{P}}{2\sqrt{s}}\frac{L}{2L+1}\sigma_{\mu\nu}^\perp \quad (45)$$

$$X_{\mu\mu_2\dots\mu_L}(k^\perp)X_{\nu\mu_2\dots\mu_L}(q^\perp)u(q_1)BW_L^+(s)$$

The formulas for the convolution of X -operators with one free index in each operator is given in Appendix B, eq.(167). Only the last, antisymmetric term, gives a nonzero result:

$$A = (\sqrt{k_\perp^2}\sqrt{q_\perp^2})^L \bar{u}(k_1)\frac{\sqrt{s} + \hat{P}}{2\sqrt{s}}\frac{\alpha(L)}{2L+1}BW_L^+(s) \left[(L+1)P_L(z) - \frac{\sigma_{\mu\nu}k_\mu q_\nu}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}}P'_L(z) \right] u(q_1) \quad (46)$$

Let us now construct the vertices for the decay of composite particles with spin-parity $1/2^+, 3/2^-, 5/2^+ \dots$ into πN . The state with $1/2^+$ is a scalar in tensor space and decays into πN with $L = 1$. Therefore this scalar should be constructed from k_μ^\perp . It cannot be $\hat{k}^\perp = k_\mu^\perp \gamma_\mu$ since such an operator is not orthogonal to the $1/2^-$ state:

$$\bar{u}(P)\hat{k}^\perp u(k_1) = \bar{u}(P)(\hat{k}_1 - \alpha\hat{P})u(k_1) = \bar{u}(P)u(k_1)(m_1 - a(s)\sqrt{s}). \quad (47)$$

Here we used:

$$k_{1\mu} = k_\mu^\perp + a(s)P_\mu, \quad (48)$$

$$\text{with } a(s) = \frac{Pk_1}{P^2} = \frac{s + m_N^2 - m_\pi^2}{2s}$$

Changing the parity in fermion sector can be done by adding a γ_5 matrix. Then the basic operator for the decay of a $1/2^+$ state into a nucleon and a pseudoscalar meson has the form:

$$i\gamma_5 \hat{k}^\perp \quad (49)$$

where \hat{k}^\perp is introduced just for convenience. Indeed

$$\bar{u}(P)i\gamma_5 \hat{k}^\perp u(k_1) = \bar{u}(P)i\gamma_5(\hat{k}_1 - a(s)\hat{P})u(k_1) = \bar{u}(P)i\gamma_5 u(k_1)(m_1 + a(s)\sqrt{s}). \quad (50)$$

Let us denote the last expression in (50) as χ :

$$\chi_i = m_i + a(s)\sqrt{s} \rightarrow (\text{in c.m.s}) m_i + k_{i0} \quad (51)$$

In general one can also introduce another scalar expression using γ matrices and k^\perp :

$$\varepsilon_{ijkl}\gamma_i\gamma_j k_k^\perp P_l \quad (52)$$

where ε_{ijkl} is the antisymmetric tensor. However using the properties of the γ matrices:

$$i\gamma^5\gamma_i\gamma_j\gamma_k = \varepsilon_{ijkl}\gamma_l \quad (53)$$

one can show that this operator is identical to (49).

For the decay of systems with $J = L - 1/2$ into πN we obtain:

$$\bar{\Psi}_{\mu_1\dots\mu_{L-1}}i\gamma_5\gamma_\nu X_{\nu\mu_1\dots\mu_{L-1}}^{(L)}(k^\perp)u(k_1) \quad (54)$$

Therefore the vertex function can be written as:

$$\bar{\Psi}_{\mu_1\dots\mu_{L-1}}N_{\mu_1\dots\mu_{L-1}}^-(k^\perp)u(k_1) \quad (55)$$

$$N_{\mu_1\dots\mu_{L-1}}^-(k^\perp) = i\gamma_5\gamma_\nu X_{\nu\mu_1\dots\mu_{L-1}}^{(L)}(k^\perp) \quad (56)$$

leading to the following amplitude for the transition $\pi N \rightarrow R \rightarrow \pi N$:

$$A = \bar{u}(k_1)X_{\alpha\mu_1\dots\mu_{L-1}}^{(L)}(k)\gamma_\alpha^\perp i\gamma_5 F_{\nu_1\dots\nu_{L-1}}^{\mu_1\dots\mu_{L-1}}(P)i\gamma_5\gamma_\xi^\perp X_{\xi\nu_1\dots\nu_{L-1}}^{(L)}(q)u(q_1)BW_L^-(s) \quad (57)$$

$$A = (\sqrt{k_\perp^2}\sqrt{q_\perp^2})^{L-1}BW_L^-(s)\bar{u}_i(k_1)\frac{\sqrt{s} + \hat{P}}{2\sqrt{s}}\frac{\alpha(L)}{2L-1} \left[\hat{k}^\perp \hat{q}^\perp L P_{L-1}(z) - \hat{k}^\perp \frac{\sigma_{\mu\nu}k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}} \hat{q}^\perp P'_{L-1}(z) \right] u_f(q_1)$$

Taking into account that

$$\hat{k}^\perp \hat{q}^\perp = (k^\perp q^\perp) + \sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp = \sqrt{k_\perp^2}\sqrt{q_\perp^2} \left(z + \frac{\sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}} \right) \quad (58)$$

$$\hat{k}^\perp \frac{\sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}} \hat{q}^\perp = \sqrt{k_\perp^2}\sqrt{q_\perp^2} \left(1 - z^2 - z \frac{\sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}} \right)$$

(Remember $z = (k^\perp q^\perp)/(\sqrt{k_\perp^2}\sqrt{q_\perp^2})$) we obtain:

$$A = (\sqrt{k_\perp^2}\sqrt{q_\perp^2})^L BW_L^-(s)\bar{u}_i(k_1)\frac{\sqrt{s} + \hat{P}}{2\sqrt{s}}\frac{\alpha(L)}{L} \times \left[(LzP_{L-1}(z) - (1 - z^2)P'_{L-1}(z)) + \frac{\sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}}(LP_{L-1} + zP'_{L-1}(z)) \right] u_f(q_1). \quad (59)$$

Using the properties of Legendre polynomials (given in Appendix A) the final expression for πN scattering due to ' \perp ' resonances reads:

$$A = (\sqrt{k_\perp^2}\sqrt{q_\perp^2})^L BW_L^-(s)\bar{u}_i(k_1)\frac{\sqrt{s} + \hat{P}}{2\sqrt{s}}\frac{\alpha(L)}{L} \times \left[LP_L(z) + \frac{\sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}}P'_L(z) \right] u_f(q_1) \quad (60)$$

Therefore the total $\pi N \rightarrow \pi N$ transition amplitude is equal to:

$$A = (\sqrt{k_\perp^2}\sqrt{q_\perp^2})^L \bar{u}_i(k_1)\frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} \left[f_1 - \frac{\sigma_{\mu\nu}^\perp k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2}\sqrt{q_\perp^2}} f_2 \right] u_f(q_1)$$

$$f_1 = \sum_L \left[\frac{\alpha(L)}{2L+1}(L+1)BW_L^+(s) + \frac{\alpha(L)}{L}L BW_L^-(s) \right] P_L(z)$$

$$f_2 = \sum_L \left[\frac{\alpha(L)}{2L+1}BW_L^+(s) - \frac{\alpha(L)}{L}BW_L^-(s) \right] P'_L(z) \quad (61)$$

Let us calculate the amplitude (61) in the c.m.s. of the resonance where $P = (\sqrt{s}, \mathbf{0})$:

$$\begin{aligned} \bar{u}_i(k_1) \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} u_f(q_1) = \\ \frac{((k_{10} + m)\omega^*, -(\mathbf{k}_1\sigma)\omega^*)}{\sqrt{k_{10} + m}} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \frac{\begin{pmatrix} (q_{10} + m)\omega' \\ (\mathbf{q}_1\sigma)\omega' \end{pmatrix}}{\sqrt{q_{10} + m}} = \\ \omega^* \sqrt{\chi_i \chi_f} \omega' \\ \sigma_{\mu\nu} = \begin{pmatrix} -i\varepsilon_{\mu\nu j} \sigma_j & 0 \\ 0 & -i\varepsilon_{\mu\nu j} \sigma_j \end{pmatrix} = -i\varepsilon_{\mu\nu j} \sigma_j I \end{aligned} \quad (62)$$

Here, ω and ω' are two-dimensional spinors of the initial and final state nucleons. Thus

$$A = (-1)^L (|\mathbf{k}||\mathbf{q}|)^L \sqrt{\chi_i \chi_f} \omega^* \left[f_1 - i\varepsilon_{\mu\nu j} \frac{\sigma_j k_\mu q_\nu}{|\mathbf{k}||\mathbf{q}|} f_2 \right] \omega'.$$

Defining the vector normal to the decay plane as

$$\mathbf{n}_j = -\varepsilon_{\mu\nu j} \frac{k_\mu q_\nu}{|\mathbf{k}||\mathbf{q}|} \quad (63)$$

we obtain the final expression

$$A = (-1)^L (|\mathbf{k}||\mathbf{q}|)^L \omega^* \sqrt{\chi_i \chi_f} [f_1 + i(\boldsymbol{\sigma}\mathbf{n})f_2] \omega \quad (64)$$

When fitting πN scattering data, the following expression (defined in the c.m.s.) is often used

$$\begin{aligned} A_{\pi N} &= \omega^* [G(s, t) + H(s, t) i(\boldsymbol{\sigma}\mathbf{n})] \omega' \\ G(s, t) &= \sum_L [(L+1)F_L^+(s) - LF_L^-(s)] P_L(z) \\ H(s, t) &= \sum_L [F_L^+(s) + F_L^-(s)] P_L'(z) \end{aligned} \quad (65)$$

The F_L^\pm are functions which depend only on energy. Comparing our expressions with (65) we obtain the following correspondence:

$$\begin{aligned} F_L^+ &= (-1)^{L+1} (|\mathbf{k}||\mathbf{q}|)^L \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{2L+1} BW_L^+(s) \\ F_L^- &= (-1)^L (|\mathbf{k}||\mathbf{q}|)^L \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{L} BW_L^-(s) \end{aligned} \quad (66)$$

5 Operators for the decay of baryons into a nucleon and a vector particle

A vector particle (e.g. a virtual photon γ^* or a ρ -meson) has spin 1 and therefore the γ^*N system can form two spin states with $S = 1/2$ and $3/2$. In combination with the orbital angular momentum, six sets of partial waves can be formed

$$\begin{aligned} J &= L_{\gamma N} + \frac{1}{2}, \quad S = \frac{1}{2}, \quad P = (-1)^{L_{\gamma N}+1}, \quad L_{\gamma N} = 0, 1, \dots \\ J &= L_{\gamma N} - \frac{3}{2}, \quad S = \frac{3}{2}, \quad P = (-1)^{L_{\gamma N}+1}, \quad L_{\gamma N} = 2, 3, \dots \\ J &= L_{\gamma N} + \frac{1}{2}, \quad S = \frac{3}{2}, \quad P = (-1)^{L_{\gamma N}+1}, \quad L_{\gamma N} = 1, 2, \dots \end{aligned}$$

$$\begin{aligned} J &= L_{\gamma N} - \frac{1}{2}, \quad S = \frac{1}{2}, \quad P = (-1)^{L_{\gamma N}+1}, \quad L_{\gamma N} = 1, 2, \dots \\ J &= L_{\gamma N} - \frac{1}{2}, \quad S = \frac{3}{2}, \quad P = (-1)^{L_{\gamma N}+1}, \quad L_{\gamma N} = 1, 2, \dots \\ J &= L_{\gamma N} + \frac{3}{2}, \quad S = \frac{3}{2}, \quad P = (-1)^{L_{\gamma N}+1}, \quad L_{\gamma N} = 0, 1, \dots \end{aligned} \quad (67)$$

5.1 Operators for $1/2^-$, $3/2^+$, $5/2^- \dots$ states

Let us start from the operators for the '+' states. A $1/2^-$ baryon decays into a baryon with $J^P = 1/2^+$ and a vector particle in either S or D-wave. In case of an S-wave decay the orbital angular momentum operator is a unit operator and the polarization vector can be convoluted only with a γ matrix. However the γ matrix changes the parity of the system. To compensate this unwanted change an additional γ_5 matrix has to be introduced. Therefore the operator describing the transition of the state with spin $1/2^-$ into a γ and $1/2^+$ fermion in S-wave is

$$\bar{u}(P) i\gamma_\mu \gamma_5 u(k_1) \varepsilon_\mu \quad (68)$$

Here $\bar{u}(P)$ is the bispinor describing a baryon resonance with momentum P , $u(k_1)$ is the bispinor for the final fermion with momentum k_1 and ε_μ is the polarization vector of the vector particle. The operator (68) is a $1/2$ spin operator and its combination with the orbital angular momentum operators $X_{\mu_1 \dots \mu_n}^{(n)}$ defines the first set of the operators (67):

$$\bar{\Psi}_{\alpha_1 \dots \alpha_L} \gamma_\mu i\gamma_5 X_{\alpha_1 \dots \alpha_L}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu \quad (69)$$

As before, $\Psi_{\alpha_1 \dots \alpha_L}$ is a fermionic bispinor wave function with spin $J = L + 1/2$, and k^\perp is the component of the relative momentum of the γ^*N system orthogonal to the total momentum of the system. For these partial waves the orbital angular momentum in the γ^*N system coincides with orbital angular momentum in πN which we denote as L .

The decay of a $1/2^-$ state into a $1/2^+$ and a vector particle in D-wave must be described by the D-wave orbital angular momentum operator:

$$\bar{u}(P) \gamma_\nu i\gamma_5 X_{\mu\nu}^{(2)}(k^\perp) u(k_1) \varepsilon_\mu \quad (70)$$

One can easily write down the whole set of such operators with $J = L_{\gamma N} - 3/2$ by

$$\bar{\Psi}_{\alpha_1 \dots \alpha_L} \gamma_\nu i\gamma_5 X_{\mu\nu\alpha_1 \dots \alpha_L}^{(L+2)}(k^\perp) u(k_1) \varepsilon_\mu \quad (71)$$

Remember that L is the orbital angular momentum in the decay of a resonance into πN ($L_{\gamma N} = L + 2$).

The third set of operators starts from the total momentum $3/2$. The basic operator describes the P-wave decay of a $3/2^+$ system into a baryon and a vector particle. It has the form

$$\bar{\Psi}_\mu \gamma_\nu i\gamma_5 X_\nu^{(1)}(k^\perp) u(k_1) \varepsilon_\mu \quad (72)$$

The operators for a baryon with $J = L_{\gamma N} + 1/2$ can be written as

$$\bar{\Psi}_{\mu\alpha_1 \dots \alpha_{L-1}} \gamma_\nu i\gamma_5 X_{\nu\alpha_1 \dots \alpha_{L-1}}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu \quad (73)$$

In case of photoproduction rather than electroproduction the operators (71) are reduced due to gauge invariance to those given in (69). Gauge invariance requires

$$\varepsilon_\mu k_{1\mu} = \varepsilon_\mu k_{2\mu} = \varepsilon_\mu k_\mu^\perp = 0 \quad (74)$$

Using (29) we obtain

$$\begin{aligned} & \bar{\Psi}_{\alpha_1 \dots \alpha_L} \gamma_\nu i \gamma_5 X_{\mu\nu\alpha_1 \dots \alpha_L}^{(L+2)}(k^\perp) u(k_1) \varepsilon_\mu = \\ & \frac{-k_\perp^2 \alpha(L)}{(2L-1)\alpha(L-2)} \bar{\Psi}_{\alpha_1 \dots \alpha_L} \gamma_\mu i \gamma_5 X_{\alpha_1 \dots \alpha_L}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu \quad (75) \end{aligned}$$

Although operators (71) applied to the case of real photons produce the same angular dependence as the operators (69), the former can provide an additional energy dependence which can be important for broad states.

It is convenient to write the decay amplitudes as a convolution of the bispinor wave functions and the vertex functions $V_{\alpha_1 \dots \alpha_L}^{(i+)\mu}$ $i = 1, 2, 3$. Then eqns.(69,71,73) can be rewritten as

$$\begin{aligned} & \bar{\Psi}_{\alpha_1 \dots \alpha_L} V_{\alpha_1 \dots \alpha_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu \\ & V_{\alpha_1 \dots \alpha_L}^{(1+)\mu}(k^\perp) = \gamma_\mu i \gamma_5 X_{\alpha_1 \dots \alpha_L}^{(L)}(k^\perp) \quad (76) \\ & V_{\alpha_1 \dots \alpha_L}^{(2+)\mu}(k^\perp) = \gamma_\nu i \gamma_5 X_{\mu\nu\alpha_1 \dots \alpha_L}^{(L+2)}(k^\perp) \\ & V_{\alpha_1 \dots \alpha_L}^{(3+)\mu}(k^\perp) = \gamma_\nu i \gamma_5 X_{\nu\alpha_1 \dots \alpha_{L-1}}^{(L)}(k^\perp) g_{\mu\alpha_L}^\perp \end{aligned}$$

In the helicity approach the property discussed above means that a $1/2$ state is described by only one helicity amplitude while states with higher spin are described by helicity $1/2$ and $3/2$ amplitudes.

5.2 Operators for $1/2^+$, $3/2^-$, $5/2^+$... states

A $1/2^+$ particle decays into a fermion with $J^P = 1/2^+$ and spin-1 particle in relative P-wave only. The operator for spin $1/2$ of the γ^*N system can be constructed in the same way as the corresponding operator for the $'+'$ -states. The P-wave orbital angular momentum operator must be convoluted with a γ -matrix. In this case, the γ_5 operator is not needed to provide the correct parity. The transition amplitude can be written as

$$\bar{u}(P) \gamma_\xi \gamma_\mu X_\xi^{(1)} u(k_1) \varepsilon_\mu \quad (77)$$

and the operator for the state with $S = 1/2$ and $J = L_{\gamma N} - 1/2$ has the form

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} \gamma_\xi \gamma_\mu X_{\xi\alpha_1 \dots \alpha_{L-1}}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu \quad (78)$$

with $L \equiv L_{\pi N} = L_{\gamma N}$.

For the 'minus' states, the operators with $S = 3/2$ and $J = L_{\gamma N} - 1/2$ have the same orbital angular momentum as the $S = 1/2$ operator. However here the polarization vector convoluted with the index of the orbital angular momentum operator. Then

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} X_{\mu\alpha_1 \dots \alpha_{L-1}}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu \quad (79)$$

The third set of operators starts from total spin $3/2$. The basic operator describes the decay of the $3/2^-$ system into the nucleon and a photon in relative S-wave. Thus

$$\bar{\Psi}_\mu u(k_1) \varepsilon_\mu, \quad (80)$$

and we obtain the set:

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} X_{\alpha_2 \dots \alpha_{L-1}}^{(L-2)}(k^\perp) g_{\alpha_1\mu}^\perp u(k_1) \varepsilon_\mu \quad (81)$$

Remember that for these states $L = L_{\gamma N} + 2$.

For real photons the operator (79) vanishes for $J = 1/2^+$; for higher states these operators provide some additional energy dependence in the partial waves (81). For convenience we introduce vertex functions $V_{\alpha_1 \dots \alpha_{L-1}}^{(i-)\mu}$ $i = 1, 2, 3$ as it was done in the case of $'+'$ states

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} V_{\mu\alpha_1 \dots \alpha_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu$$

$$\begin{aligned} & V_{\alpha_1 \dots \alpha_{L-1}}^{(1-)\mu}(k^\perp) = \gamma_\xi \gamma_\mu X_{\xi\alpha_1 \dots \alpha_{L-1}}^{(L)}(k^\perp) \quad (82) \\ & V_{\alpha_1 \dots \alpha_{L-1}}^{(2-)\mu}(k^\perp) = X_{\mu\alpha_1 \dots \alpha_{L-1}}^{(L)}(k^\perp) \\ & V_{\alpha_1 \dots \alpha_{L-1}}^{(3-)\mu}(k^\perp) = X_{\alpha_2 \dots \alpha_{L-1}}^{(L-2)}(k^\perp) g_{\alpha_1\mu}^\perp \end{aligned}$$

6 Single meson photoproduction

The amplitude for the photoproduction of a single pseudoscalar meson (for the sake of simplicity let us take the pion) is well known and can be found in the literature (see for example [21] and references therein). The general structure of the amplitude is

$$A = \omega^* J_\mu \varepsilon_\mu \omega'$$

$$J_\mu = i f_1 \sigma_\mu + f_2(\sigma \mathbf{q}) \frac{\varepsilon_{\mu ij} \sigma_i k_j}{|\mathbf{k}| |\mathbf{q}|} + i f_3 \frac{(\sigma \mathbf{k})}{|\mathbf{k}| |\mathbf{q}|} q_\mu + i f_4 \frac{(\sigma \mathbf{q})}{\mathbf{q}^2} q_\mu \quad (83)$$

where \mathbf{q} is the momentum of the nucleon in the πN channel and \mathbf{k} is the momentum of the nucleon in the γN channel calculated in the c.m.s. of the reaction and σ_i are Pauli matrices.

The functions f_i have the following angular dependence:

$$\begin{aligned} & f_1(z) = \\ & \sum_{L=0}^{\infty} [L M_L^+ + E_L^+] P'_{L+1}(z) + [(L+1) M_L^- + E_L^-] P'_{L-1}(z) \\ & f_2(z) = \sum_{L=1}^{\infty} [(L+1) M_L^+ + L M_L^-] P'_L(z) \\ & f_3(z) = \sum_{L=1}^{\infty} [E_L^+ - M_L^+] P''_{L+1}(z) + [E_L^- + M_L^-] P''_{L-1}(z) \\ & f_4(z) = \sum_{L=2}^{\infty} [M_L^+ - E_L^+ - M_L^- - E_L^-] P''_L(z). \quad (84) \end{aligned}$$

Here L corresponds to the orbital angular momentum in the πN system, $P_L(z)$ are Legendre polynomials $z = (\mathbf{k}\mathbf{q})/(|\mathbf{k}||\mathbf{q}|)$ and E_L^\pm and M_L^\pm are electric and magnetic multipoles describing transitions to states with $J = L \pm 1/2$. In the following we will construct the $\gamma N \rightarrow \pi N$ transition amplitudes using the operators defined in the previous sections and show that in c.m.s. these amplitudes satisfy the equations (83, 84).

6.1 Photoproduction amplitudes for $1/2^-, 3/2^+, 5/2^- \dots$ states

The angular dependence of the single-meson-production amplitude via an intermediate resonance has the general form

$$\bar{u}(q_1) \tilde{N}_{\alpha_1 \dots \alpha_n}^\pm(q^\perp) F_{\beta_1 \dots \beta_n}^{\alpha_1 \dots \alpha_n}(P) V_{\beta_1 \dots \beta_n}^{(i\pm)\mu}(k^\perp) u(k_1) \varepsilon_\mu \quad (85)$$

Here q_1 and k_1 are the momenta of the nucleon in the πN and γN channel and q^\perp and k^\perp are the components of the relative momenta which are orthogonal to the total momentum of the resonance.

If states with $J = L + 1/2$ are produced from a γN partial wave with spin $1/2$ one has the following expression for the amplitude:

$$A^+(1/2) = \bar{u}(q_1) X_{\alpha_1 \dots \alpha_L}^{(L)}(q^\perp) F_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) \gamma_\mu i \gamma_5 X_{\beta_1 \dots \beta_L}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu BW(s) \quad (86)$$

where $BW(s)$ represent the dynamical part of the amplitude. Taking into account the properties of the projection operator this expression can be rewritten as

$$\begin{aligned} & \bar{u}(q_1) X_{\alpha_1 \dots \alpha_L}^{(L)}(q^\perp) T_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L} \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} X_{\beta_1 \dots \beta_L}^{(L)}(k^\perp) \\ & \gamma_\mu i \gamma_5 u(k_1) \varepsilon_\mu = \\ & \bar{u}(q_1) \left[\frac{L+1}{2L+1} X_{\alpha_1 \dots \alpha_L}^{(L)}(q^\perp) X_{\alpha_1 \dots \alpha_L}^{(L)}(k^\perp) - \right. \\ & \left. \frac{L}{2L+1} \sigma_{\alpha\beta} X_{\alpha\alpha_2 \dots \alpha_L}^{(L)}(q^\perp) X_{\beta\alpha_2 \dots \alpha_L}^{(L)}(k^\perp) \right] \\ & \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} \gamma_\mu i \gamma_5 u(k_1) \varepsilon_\mu \quad (87) \end{aligned}$$

Using the expression for the convolution of two X -operators with two external indices (as given in Appendix B) one obtains

$$\begin{aligned} A^+(1/2) &= \bar{u}(q_1) \frac{L+1}{2L+1} \alpha(L) (\sqrt{q^\perp} \sqrt{k^\perp})^L \\ & \left[P_L(z) - \frac{P'_L(z)}{L+1} \sigma_{\alpha\beta} \frac{q_\alpha^\perp k_\beta^\perp}{(\sqrt{q^\perp} \sqrt{k^\perp})} \right] \\ & \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} \gamma_\mu i \gamma_5 u(k_1) \varepsilon_\mu BW(s) \quad (88) \end{aligned}$$

In the c.m.s.

$$\bar{u}(q_1) \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} \gamma_\mu i \gamma_5 u(k_1) \varepsilon_\mu = -\sqrt{\chi_i \chi_f} i \omega^* (\varepsilon_i \sigma_i) \omega' \quad (89)$$

holds, leading to

$$\begin{aligned} A^+(1/2) &= \omega^* \sqrt{\chi_i \chi_f} \frac{L+1}{2L+1} \alpha(L) i(-\varepsilon_i) (\sqrt{q^\perp} \sqrt{k^\perp})^L \\ & \left[\sigma_i P_L(z) + i \frac{P'_L(z)}{L+1} \varepsilon_{\alpha\beta\xi} \sigma_\xi \sigma_i \frac{q_\alpha^\perp k_\beta^\perp}{(\sqrt{q^\perp} \sqrt{k^\perp})} \right] \omega' BW(s) \quad (90) \end{aligned}$$

Here all vectors are three-dimensional. Using in addition the properties of Pauli matrices

$$\sigma_i \sigma_j = \delta_{ij} + i \varepsilon_{ijk} \sigma_k \quad (91)$$

one obtains the final expression

$$\begin{aligned} A^+(1/2) &= -\omega^* \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{2L+1} \varepsilon_i (\sqrt{q^\perp} \sqrt{k^\perp})^L \\ & \left[i \sigma_i \left((L+1) P_L(z) + z P'_L(z) \right) + (\sigma \mathbf{q}) \frac{\varepsilon_{ijm} \sigma_j k_m}{|\mathbf{k}||\mathbf{q}|} P'_L(z) \right] \\ & \omega' BW(s) \quad (92) \end{aligned}$$

Taking into account the properties of the Legendre polynomials (given in Appendix A) the amplitude can be compared with equations (83), (84). One finds the following correspondence between the spin operators and multipoles:

$$\begin{aligned} E_L^{+(1)} &= (-1)^L \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{2L+1} \frac{(|\mathbf{k}||\mathbf{q}|)^L}{L+1} BW(s) \\ M_L^{+(1)} &= E_L^{+(1)} \quad (93) \end{aligned}$$

In the case of photoproduction, only two γN operators are independent for every resonance with spin $3/2$ and higher (for $J = 1/2$ states there is only one independent operator). For the set of $J = L + 1/2$ states the second operator has the amplitude structure

$$\begin{aligned} A^+(3/2) &= \bar{u}(q_1) X_{\alpha_1 \dots \alpha_L}^{(L)}(q^\perp) F_{\mu\beta_2 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) \\ & \gamma_\xi i \gamma_5 X_{\xi\beta_2 \dots \beta_L}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu BW(s) \quad (94) \end{aligned}$$

Using expressions given in Appendix B one obtains the multipole decomposition

$$\begin{aligned} E_L^{+(2)} &= (-1)^L \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{2L+1} \frac{(|\mathbf{k}||\mathbf{q}|)^L}{L+1} BW(s) \\ M_L^{+(2)} &= -\frac{E_L^{+(2)}}{L} \quad (95) \end{aligned}$$

6.2 Photoproduction amplitudes for $1/2^+, 3/2^-, 5/2^+ \dots$ states

The $\gamma N \rightarrow \pi N$ amplitude for states with $J = L - 1/2$ in the πN channel has the structure

$$\begin{aligned} A^-(1/2) &= \bar{u}(q_1) \gamma_\xi i \gamma_5 X_{\xi\alpha_1 \dots \alpha_{L-1}}^{(L)}(q^\perp) F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) \\ & \gamma_\xi \gamma_\mu X_{\xi\beta_1 \dots \beta_{L-1}}^{(L)}(k^\perp) u(k_1) \varepsilon_\mu BW(s) \quad (96) \end{aligned}$$

For amplitude (96) we find the following correspondence to the multipole decomposition (see Appendix B for details):

$$\begin{aligned} E_L^{-(1)} &= (-1)^L \sqrt{\chi_i \chi_f} |\mathbf{k}|^L |\mathbf{q}|^L \frac{\alpha(L)}{L^2} BW(s) \\ M_L^{-(1)} &= -E_L^{-(1)} \end{aligned} \quad (97)$$

Amplitudes including spin 3/2 operators have the structure

$$\begin{aligned} A^-(3/2) &= \bar{u}(q_1) \gamma_\xi i \gamma_5 X_{\xi \alpha_1 \dots \alpha_{L-1}}^{(L)}(q^\perp) F_{\mu \beta_2 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) \\ X_{\beta_2 \dots \beta_{L-1}}^{(L-2)}(k^\perp) u(k_1) \varepsilon_\mu BW(s) \end{aligned} \quad (98)$$

Using expressions in Appendix B the decomposition of this amplitude into the multipole representation is the following:

$$\begin{aligned} E_L^{-(2)} &= (-1)^L \sqrt{\chi_i \chi_f} |\mathbf{k}|^{L-2} |\mathbf{q}|^L \frac{\alpha(L-2)}{(L-1)L} BW^-(s) \\ M_L^{-(2)} &= 0 \end{aligned} \quad (99)$$

6.3 Relations between the amplitudes in the spin-orbit and helicity representation

The helicity amplitudes $A^{1/2}$ and $A^{3/2}$ are of course different from the spin 1/2 and 3/2 amplitudes $A^\pm(1/2)$, $A^\pm(3/2)$. For '+' multipoles the relations between the helicity amplitudes and multipoles are

$$\begin{aligned} A^{1/2} &= -\frac{1}{2}(LM_L^+ + (L+2)E_L^+) \\ A^{3/2} &= \frac{1}{2}\sqrt{L(L+2)}(E_L^+ - M_L^+) \end{aligned} \quad (100)$$

For the '-' sector the relations are

$$\begin{aligned} A^{1/2} &= \frac{1}{2}((L+1)M_L^- - (L-1)E_L^-) \\ A^{3/2} &= -\frac{1}{2}\sqrt{(L-1)(L+1)}(E_L^- + M_L^+) \end{aligned} \quad (101)$$

For '+' states we obtain the following decomposition of the spin 1/2 amplitude (93):

$$\begin{aligned} A^{1/2} &= -(L+1)E_L^{+(1)} \\ A^{3/2} &= 0 \end{aligned} \quad (102)$$

Obviously the spin 1/2 state can not have a helicity 3/2 projection. For the spin 3/2 state one gets

$$\begin{aligned} A^{1/2} &= -\frac{L+1}{2}E_L^{+(2)} \\ A^{3/2} &= \frac{1}{2}\sqrt{\frac{L+2}{L}}(L+1)E_L^{+(2)} \end{aligned} \quad (103)$$

The ratio of the helicity amplitudes can be calculated directly if the ratio of the spin amplitudes is known. The $BW^+(s)$ in both amplitudes is an energy dependent part

of the amplitude which depends on the model used in the analysis. If a resonance is produced and decays with radius r the regularization of the amplitude can be done with, e.g., Blatt-Weisskopf formfactors (see Appendix C). If we also explicitly extract the initial coupling constants $g_{1/2}$ and $g_{3/2}$ for the spin 1/2 and 3/2, then the expression for the total amplitude for '+' states has the form

$$\begin{aligned} A_{tot}^{L+} &= \\ [g_{1/2} A^+(1/2) + g_{3/2} A^+(3/2)] &\frac{BW(s)}{F(L, q_\perp^2, r) F(L, k_\perp^2, r)}. \end{aligned} \quad (104)$$

In this case the multipole amplitudes can be rewritten as following:

$$\begin{aligned} E_L^{+(1)} &= \\ (-1)^L \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{2L+1} \frac{(|\mathbf{k}||\mathbf{q}|)^L}{L+1} &\frac{BW(s)}{F(L, q_\perp^2, r) F(L, k_\perp^2, r)} g_{1/2} \end{aligned} \quad (105)$$

$$\begin{aligned} E_L^{+(2)} &= \\ (-1)^L \sqrt{\chi_i \chi_f} \frac{\alpha(L)}{2L+1} \frac{(|\mathbf{k}||\mathbf{q}|)^L}{L+1} &\frac{BW(s)}{F(L, q_\perp^2, r) F(L, k_\perp^2, r)} g_{3/2} \end{aligned} \quad (106)$$

From (102) and (103) one can calculate the the ratio between helicity amplitudes for '+' states:

$$\frac{A^{3/2}}{A^{1/2}} = -\frac{\frac{1}{2}\sqrt{\frac{L+2}{L}}(L+1)E_L^{+(2)}}{\frac{L+1}{2}E_L^{+(2)} + (L+1)E_L^{+(1)}} = -\sqrt{\frac{L+2}{L}} \frac{1}{1+2R}$$

This ratio does not depend on the final state of the photoproduction process and is valid for any photoproduction reaction.

In the case of the '-' states we get for the spin 1/2 amplitude:

$$\begin{aligned} A^{1/2} &= -LE_L^{-(1)} \\ A^{3/2} &= 0 \end{aligned} \quad (108)$$

and for the spin 3/2 amplitudes

$$\begin{aligned} A^{1/2} &= -\frac{L-1}{2}E_L^{-(2)} \\ A^{3/2} &= -\frac{1}{2}\sqrt{(L-1)(L+1)}E_L^{-(2)} \end{aligned} \quad (109)$$

Here the production vertices have the different orbital angular momenta and therefore:

$$A_{tot}^{L-} = \left[g_{1/2} \frac{A^-(1/2)}{F(L, k_\perp^2, r)} + g_{3/2} \frac{A^-(3/2)}{F(L-2, k_\perp^2, r)} \right] \frac{BW(s)}{F(L, q_\perp^2, r)}$$

The multipole amplitudes can be rewritten as follows:

$$\begin{aligned} E_L^{-(1)} &= \\ (-1)^L \sqrt{\chi_i \chi_f} |\mathbf{k}|^L |\mathbf{q}|^L \frac{\alpha(L)}{L^2} &\frac{BW(s)}{F(L, q_\perp^2, r) F(L, k_\perp^2, r)} g_{1/2} \end{aligned} \quad (110)$$

$$E_L^{-(2)} = (-1)^L \sqrt{\chi_i \chi_f} |\mathbf{k}|^{L-2} |\mathbf{q}|^L \frac{\alpha(L-2)}{(L-1)L} \frac{BW(s)}{F(L, q_\perp^2, r) F(L-2, k_\perp^2, r)} g_{3/2} \quad (111)$$

For the ratio of helicity amplitudes one obtains:

$$\frac{A^{3/2}}{A^{1/2}} = \frac{\frac{1}{2} \sqrt{(L-1)(L+1)} E_L^{-(2)}}{\frac{L-1}{2} E_L^{-(2)} + L E_L^{-(1)}} = \sqrt{\frac{L+1}{L-1}} \frac{1}{1+2R\kappa} \quad (112)$$

where

$$\kappa = \frac{(2L-1)(2L-3)}{L(L-1)} |\mathbf{k}|^2 \frac{F(L-2, k_\perp^2, r)}{F(L, k_\perp^2, r)} \quad (113)$$

7 Operators for the decay of a baryon resonance into pseudoscalar meson and a baryon with spin 3/2

The spin of a pseudoscalar meson and of a baryon with spin 3/2 combine to a total spin 3/2. In combination with the orbital angular momentum operators four sets of partial waves can be formed:

$$\begin{aligned} J &= L_\Delta - 3/2, \quad S = 3/2, \quad P = (-1)^{L_\Delta+1}, \quad L_\Delta = 2, 3, \dots \\ J &= L_\Delta + 1/2, \quad S = 3/2, \quad P = (-1)^{L_\Delta+1}, \quad L_\Delta = 1, 2, \dots \\ J &= L_\Delta - 1/2, \quad S = 3/2, \quad P = (-1)^{L_\Delta+1}, \quad L_\Delta = 1, 2, \dots \\ J &= L_\Delta + 3/2, \quad S = 3/2, \quad P = (-1)^{L_\Delta+1}, \quad L_\Delta = 0, 1, \dots \end{aligned} \quad (114)$$

7.1 Operators for $1/2^-$, $3/2^+$, $5/2^- \dots$ states

A $1/2^-$ particle decays into a $J^P = 3/2^+$ -particle and pseudoscalar meson in D-wave. Only one of the indices of the orbital angular momentum operator can be absorbed by a γ -matrix. Again, to compensate the change of parity due to the γ -matrix one has to introduce an additional γ_5 -matrix. The operator describing the transition of a state with spin $1/2^-$ into a 0^- and a $3/2^+$ state is

$$\bar{u}(P) i\gamma_5 \gamma_\nu X_{\mu\nu}^{(2)} \Psi_\mu^\Delta \quad (115)$$

where \bar{u}_p is a bispinor describing an initial state and Ψ_μ^Δ is a vector bispinor for the final spin-3/2 fermion. The first set of operators (114) derived from eq.(115) reads

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L_\Delta-2}} i\gamma_5 \gamma_\nu X_{\mu\nu \alpha_1 \dots \alpha_{L_\Delta-2}}^{(L_\Delta)} \Psi_\mu^\Delta \quad L_\Delta = 2, 3, \dots \quad (116)$$

However it is again convenient to rewrite this expression using the orbital angular momentum L . In this case $L_\Delta = L+2$, and

$$\bar{\Psi}_{\alpha_1 \dots \alpha_L} i\gamma_5 \gamma_\nu X_{\mu\nu \alpha_1 \dots \alpha_L}^{(L+2)} \Psi_\mu^\Delta \quad L = 0, 1, \dots \quad (117)$$

The second set of operators starts from total spin 3/2. The basic operator describes the decay of the $3/2^+$ system into Δ and pion in a P-wave. It has the form

$$\bar{\Psi}_\alpha i\gamma_5 \gamma_\nu X_\nu^{(1)} g_{\alpha\mu}^\perp \Psi_\mu^\Delta \quad (118)$$

The second set of the operators (114) can be written as (here $L_\Delta = L$)

$$\bar{\Psi}_{\alpha_1 \dots \alpha_L} i\gamma_5 \gamma_\nu X_{\nu \alpha_2 \dots \alpha_L}^{(L)} g_{\alpha_1 \mu}^\perp \Psi_\mu^\Delta \quad L = 1, 2, \dots \quad (119)$$

Thus the vertex functions for '+' states are

$$\begin{aligned} \bar{\Psi}_{\alpha_1 \dots \alpha_L} N_{\alpha_1 \dots \alpha_L}^{(i+)\mu} \Psi_\mu^\Delta & \quad N_{\alpha_1 \dots \alpha_L}^{(1+)\mu} = i\gamma_5 \gamma_\nu X_{\mu\nu \alpha_1 \dots \alpha_L}^{(L+2)} \\ & \quad N_{\alpha_1 \dots \alpha_L}^{(2+)\mu} = i\gamma_5 \gamma_\nu X_{\nu \alpha_2 \dots \alpha_L}^{(L)} g_{\alpha_1 \mu}^\perp \end{aligned} \quad (120)$$

7.2 Operators for $1/2^+$, $3/2^-$, $5/2^+ \dots$ states

A $1/2^+$ particle may decay into a $J^P = 3/2^+$ baryon and 0^- meson in P-wave. In this case the P-wave orbital angular momentum operator must be converted with the vector bispinor Ψ_μ^Δ . The γ_5 operator is not needed to provide a correct parity for the state. Then

$$\bar{u}(P) X_\mu^{(1)} \Psi_\mu^\Delta \quad (121)$$

The operator for the state with $S = 3/2$ and $J = L - 1/2$ ($L = L_\Delta$) has the form

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} X_{\mu \alpha_1 \dots \alpha_{L-1}}^{(L)} \Psi_\mu^\Delta \quad L = 1, 2, \dots \quad (122)$$

As before, the second set of operators starts from total spin $S = 3/2$. The basic operator describes the decay of the $3/2^-$ system into a $3/2^+$ particle and pion in S-wave. Thus

$$\bar{\Psi}_\mu \Psi_\mu^\Delta, \quad (123)$$

and we obtain for this set

$$\bar{\Psi}_{\mu \alpha_1 \dots \alpha_{L_\Delta}} X_{\alpha_1 \dots \alpha_{L_\Delta}}^{(L_\Delta)} \Psi_\mu^\Delta \quad L_\Delta = 0, 1, \dots \quad (124)$$

Here $L = L_\Delta + 2$ and the amplitude can be rewritten as

$$\bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} X_{\alpha_2 \dots \alpha_{L-1}}^{(L-2)} g_{\alpha_1 \mu}^\perp \Psi_\mu^\Delta \quad L = 2, 3, \dots \quad (125)$$

The vertex functions for '-' states are given by:

$$\begin{aligned} \bar{\Psi}_{\alpha_1 \dots \alpha_{L-1}} N_{\alpha_1 \dots \alpha_{L-1}}^{(i-)\mu} \Psi_\mu^\Delta & \quad N_{\alpha_1 \dots \alpha_{L-1}}^{(1-)\mu} = X_{\mu \alpha_1 \dots \alpha_{L-1}}^{(L)} \\ & \quad N_{\alpha_1 \dots \alpha_{L-1}}^{(2-)\mu} = X_{\alpha_2 \dots \alpha_{L-1}}^{(L-2)} g_{\alpha_1 \mu}^\perp \end{aligned} \quad (126)$$

7.3 Operators for the decay into states with different parity

The operators given in the previous sections provide a full set of operators for the decay of a baryon into meson with spin 0 and fermion with spin 3/2. Indeed, for construction of operators only the total spin of the system plays the role. Thus the operators for $J^+ \rightarrow 0^- + 3/2^+$ decays have the same form as the operators for $J^+ \rightarrow 0^+ + 3/2^-$, $J^- \rightarrow 0^+ + 3/2^+$ and $J^- \rightarrow 0^- + 3/2^-$ decays.

8 Double pion photoproduction amplitudes

Let us construct the amplitudes for double pion photoproduction. Here reactions as shown in Fig. 1 are taken into account where the decay into the final state proceeds via production an intermediate baryon or meson resonance. The general form of the angular dependent part of the




Fig. 1. Photoproduction of two mesons due to the cascade of a resonance

amplitude for such a process is

$$\begin{aligned} & \bar{u}(q_1) \tilde{N}_{\alpha_1 \dots \alpha_n}^+(R_2 \rightarrow \mu N) F_{\beta_1 \dots \beta_n}^{\alpha_1 \dots \alpha_n}(q_1 + q_2) \\ & \tilde{N}_{\gamma_1 \dots \gamma_m}^{(j)\beta_1 \dots \beta_n}(R_1 \rightarrow \mu R_2) F_{\xi_1 \dots \xi_m}^{\gamma_1 \dots \gamma_m}(P) V_{\xi_1 \dots \xi_m}^{(i)\mu}(R_1 \rightarrow \gamma N) \\ & u(k_1) \varepsilon_\mu, \quad P = q_1 + q_2 + q_3 = k_1 + k_2 \quad (127) \end{aligned}$$

The resonance R_1 with spin $J = m + 1/2$ is produced in γN interaction, propagates and then decays into a meson and a baryon resonance R_2 with spin $J = n + 1/2$. Then the resonance R_2 propagates and decays into the final meson and a nucleon.

In the following the full vertex functions used for the construction of amplitudes are given here for convenience of the reader. One should remember that the \tilde{N} functions are different from N -functions by the order of γ -matrices. For $R \rightarrow 0^- + 1/2^+$ transitions

$$\tilde{N}_{\mu_1 \dots \mu_n}^+ = X_{\mu_1 \dots \mu_n}^{(n)} \quad \tilde{N}_{\mu_1 \dots \mu_n}^- = i\gamma_\nu \gamma_5 X_{\nu \mu_1 \dots \mu_n}^{(n+1)} \quad (128)$$

holds, while we have

$$\begin{aligned} \tilde{N}_{\alpha_1 \dots \alpha_n}^{(1+)\mu} &= i\gamma_\nu \gamma_5 X_{\mu \nu \alpha_1 \dots \alpha_n}^{(n+2)} & \tilde{N}_{\alpha_1 \dots \alpha_n}^{(1-)\mu} &= X_{\mu \alpha_1 \dots \alpha_n}^{(n+1)} \\ \tilde{N}_{\alpha_1 \dots \alpha_n}^{(2+)\mu} &= i\gamma_\nu \gamma_5 X_{\nu \alpha_2 \dots \alpha_n}^{(n)} g_{\alpha_1 \mu}^\perp & \tilde{N}_{\alpha_1 \dots \alpha_n}^{(2-)\mu} &= X_{\alpha_2 \dots \alpha_n}^{(n-1)} g_{\alpha_1 \mu}^\perp \end{aligned} \quad (129)$$

for $R \rightarrow 0^- + 3/2^+$ transitions, and

$$\begin{aligned} V_{\alpha_1 \dots \alpha_n}^{(1+)\mu} &= \gamma_\mu i\gamma_5 X_{\alpha_1 \dots \alpha_n}^{(n)} & V_{\alpha_1 \dots \alpha_n}^{(1-)\mu} &= \gamma_\xi \gamma_\mu X_{\xi \alpha_1 \dots \alpha_n}^{(n+1)} \\ V_{\alpha_1 \dots \alpha_n}^{(2+)\mu} &= \gamma_\nu i\gamma_5 X_{\mu \nu \alpha_1 \dots \alpha_n}^{(n+2)} & V_{\alpha_1 \dots \alpha_n}^{(2-)\mu} &= X_{\mu \alpha_1 \dots \alpha_n}^{(n+1)} \\ V_{\alpha_1 \dots \alpha_n}^{(3+)\mu} &= \gamma_\nu i\gamma_5 X_{\nu \alpha_1 \dots \alpha_n}^{(n+1)} g_{\mu \alpha_n}^\perp & V_{\alpha_1 \dots \alpha_n}^{(3-)\mu} &= X_{\alpha_2 \dots \alpha_n}^{(n-1)} g_{\alpha_1 \mu}^\perp \end{aligned} \quad (130)$$

for $R \rightarrow 1^- + 1/2^+$ transitions. Here n is related to the total spin of the resonance by $J = n + 1/2$.

8.1 Amplitudes for baryons states decaying into a $1/2$ state and a pion

In this section explicit expressions for the angular dependent part of the amplitudes are given for the case of a baryon produced in a $\gamma^* N$ collision. The baryon decays into a pseudoscalar particle and another (intermediate) baryon with spin $1/2$ (decaying in turn into meson and nucleon).

8.1.1 The $1/2^-$, $3/2^+$, $5/2^- \dots$ states

The amplitude for a $'+$ ' state (R_1) produced in a $\gamma^* N$ collision in a partial wave (i) decaying into a 0^- meson and an intermediate $1/2^+$ baryon (R_2) has the form

$$\begin{aligned} A^{(i)} &= \bar{u}(q_1) \tilde{N}^-(q_{12}^\perp) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} \tilde{N}_{\alpha_1 \dots \alpha_L}^+(q_1^\perp) \\ & F_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) V_{\beta_1 \dots \beta_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu \\ &= \bar{u}(q_1) i\hat{q}_{12}^\perp \gamma_5 \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} X_{\alpha_1 \dots \alpha_L}^{(L)}(q_1^\perp) \frac{\sqrt{s} + \hat{P}}{2\sqrt{s}} \\ & R_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L} V_{\beta_1 \dots \beta_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu \quad (131) \end{aligned}$$

where the k_1 and q_1 are the momenta of the nucleon in the initial and the final state, $k^\perp = 1/2(k_1 - k_2)^\perp$ and $q_1^\perp = 1/2(q_1 + q_2 - q_3)^\perp$ are their components orthogonal to the total momentum of the first resonance R_1 . Further, $s_{12} = (q_1 + q_2)^2$ and the factors $1/(2\sqrt{s})$ and $1/(2\sqrt{s_{12}})$ are introduced to suppress the divergency of the numerator of the fermion propagators at large energies. The relative momentum q_{12}^\perp is the component of q_1 and q_2 orthogonal to the total momentum $q_1 + q_2$. It is given by:

$$q_{12}^\perp = \frac{1}{2}(q_1 - q_2)_\nu \left(g_{\mu\nu} - \frac{(q_1 + q_2)_\mu (q_1 + q_2)_\nu}{(q_1 + q_2)^2} \right) \quad (132)$$

The vertex functions (128)-(129) are given for the case when the nucleon wave function is placed on the right-hand side of the amplitude. Therefore the order of the γ -matrices needs to be changed for the meson-nucleon vertices in eq.(127).

If the baryon R_2 has spin $1/2^-$ one has to construct the vertex for decay of $'+$ ' states into a 0^- and a $1/2^-$ particle. However such operators coincide with the operators for the decay of $'-$ ' states into a $0^- + 1/2^+$ system. Therefore

$$\begin{aligned} A^{(i)} &= \bar{u}(q_1) \tilde{N}^+(q_{12}^\perp) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} \tilde{N}_{\alpha_1 \dots \alpha_L}^-(q_1^\perp) \\ & F_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) V_{\beta_1 \dots \beta_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu = \\ & \bar{u}(q_1) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} i\gamma_\nu \gamma_5 X_{\nu \alpha_1 \dots \alpha_L}^{(L+1)}(q_1^\perp) \\ & F_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) V_{\beta_1 \dots \beta_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu \quad (133) \end{aligned}$$

In case of the photoproduction with real photons, the $V_{\beta_1 \dots \beta_L}^{(2+)\mu}$ vertex is reduced to $V_{\beta_1 \dots \beta_L}^{(1+)\mu}$, and can be omitted.

8.1.2 The $1/2^+$, $3/2^-$, $5/2^+$... states

If a $\prime\prime$ state is produced in a γ^*N interaction and then decays into a pseudoscalar pion and $1/2^+$ baryon the amplitude has the structure

$$\begin{aligned} A^{(i)} &= \bar{u}(q_1) \tilde{N}^-(q_{12}^\perp) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} \tilde{N}_{\alpha_1 \dots \alpha_{L-1}}^-(q_1^\perp) \\ &F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) V_{\beta_1 \dots \beta_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu = \\ &\bar{u}(q_1) i \hat{q}_{12}^\perp \gamma_5 \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} i \gamma_\nu \gamma_5 X_{\nu \alpha_1 \dots \alpha_{L-1}}^{(L)}(q_1^\perp) \\ &F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) V_{\beta_1 \dots \beta_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu \end{aligned} \quad (134)$$

If the intermediate baryon has spin $1/2^-$ then:

$$\begin{aligned} A^{(i)} &= \bar{u}(q_1) \tilde{N}^+(q_{12}^\perp) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} \tilde{N}_{\alpha_1 \dots \alpha_{L-1}}^+(q_1^\perp) \\ &F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) V_{\beta_1 \dots \beta_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu = \\ &\bar{u}(q_1) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} X_{\alpha_1 \dots \alpha_L}^{(L)}(q_1^\perp) \\ &F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) V_{\beta_1 \dots \beta_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu \end{aligned} \quad (135)$$

For photoproduction with real photons only amplitudes with $V^{(1-)}$ and $V^{(3-)}$ vertex functions should be taken into account.

8.2 Photoproduction amplitudes for baryon states decaying into a $3/2$ state and a pseudoscalar meson

Experimentally important is photoproduction of resonances decaying into $\Delta(1232)\pi$ followed by a $\Delta(1232)$ decay into a nucleon and a pion.

8.2.1 The $1/2^-$, $3/2^+$, $5/2^-$... states decaying into a meson with spin 0 and a baryon with spin $3/2$

The $\prime\prime$ states produced in a γ^*N collision can decay into a pseudoscalar meson and an intermediate baryon with spin $3/2^+$ in two partial waves. The amplitude depends on indices (ij) where index (i) is related, as before, to the partial wave in the γN channel while index (j) is related to the partial wave in the decay of the resonance into the spin 0 meson and the $3/2$ resonance R_2 .

$$\begin{aligned} A^{(ij)} &= \bar{u}(q_1) \tilde{N}_\delta^+(q_{12}^\perp) F_\nu^\delta(q_1 + q_2) \tilde{N}_{\alpha_1 \dots \alpha_L}^{(j+)\nu}(q_1^\perp) \\ &F_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) V_{\beta_1 \dots \beta_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu \end{aligned} \quad (136)$$

If the intermediate baryon R_2 has $J^P = 3/2^-$, the structure of the amplitude structure is

$$\begin{aligned} A^{(ij)} &= \bar{u}(q_1) \tilde{N}_\delta^-(q_{12}^\perp) F_\nu^\delta(q_1 + q_2) \tilde{N}_{\alpha_1 \dots \alpha_L}^{(j-)\nu}(q_1^\perp) \\ &F_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L}(P) V_{\beta_1 \dots \beta_L}^{(i+)\mu}(k^\perp) u(k_1) \varepsilon_\mu \end{aligned} \quad (137)$$

8.2.2 $1/2^+$, $3/2^-$, $5/2^+$... states decaying into a 0^- meson and a $3/2^+$ baryon

The amplitudes for $\prime\prime$ states decaying into 0^- meson and $3/2^+$ intermediate baryon are

$$\begin{aligned} A^{(ij)} &= \bar{u}(q_1) \tilde{N}_\delta^+(q_{12}^\perp) F_\nu^\delta(q_1 + q_2) \tilde{N}_{\alpha_1 \dots \alpha_{L-1}}^{(j-)\nu}(q_1^\perp) \\ &F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) V_{\beta_1 \dots \beta_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu \end{aligned} \quad (138)$$

and if the intermediate baryon R_2 has the quantum numbers $3/2^-$

$$\begin{aligned} A^{(ij)} &= \bar{u}(q_1) \tilde{N}_\delta^-(q_{12}^\perp) F_\nu^\delta(q_1 + q_2) \tilde{N}_{\alpha_1 \dots \alpha_{L-1}}^{(j+)\nu}(q_1^\perp) \\ &F_{\beta_1 \dots \beta_{L-1}}^{\alpha_1 \dots \alpha_{L-1}}(P) V_{\beta_1 \dots \beta_{L-1}}^{(i-)\mu}(k^\perp) u(k_1) \varepsilon_\mu \end{aligned} \quad (139)$$

9 t- and u-channel exchange amplitudes

Meson exchange in the t-channel plays an important role in both, in photoproduction and in pion induced reactions. Especially at large energies this mechanism dominates. In the resonance region we expect that production of baryon resonances in the s-channel dominates the interaction, at least when neutral mesons are produced. Nevertheless the t- and u-channel exchanges must be taken into account carefully.

The most straight-forward parameterization of particle exchange amplitudes is the exchange of Regge trajectories. For construction of a cross-symmetrical amplitude it is convenient to use the variable

$$\nu = \frac{1}{2}(s - u).$$

The amplitude for t-channel exchange can be written as

$$A = g_1(t) g_2(t) \frac{1 + \xi \exp(-i\pi\alpha(t))}{\sin(\pi\alpha(t))} \left(\frac{\nu}{\nu_0} \right)^{\alpha(t)} \quad (140)$$

Here g_i are vertex functions, $\alpha(t)$ is the function which describes the trajectory, ν_0 is a normalization factor (which can be taken to be 1) and ξ is the signature of the trajectory. The Pomeron, f_0 and π exchanges have a positive signature while ρ , ω and a_1 exchanges have a negative one.

Accordingly, the Reggeon propagators can be written as

$$\begin{aligned} R(+, \nu, t) &= \frac{e^{-i\frac{\pi}{2}\alpha(t)}}{\sin(\frac{\pi}{2}\alpha(t))} \left(\frac{\nu}{\nu_0} \right)^{\alpha(t)}, \\ R(-, \nu, t) &= \frac{i e^{-i\frac{\pi}{2}\alpha(t)}}{\cos(\frac{\pi}{2}\alpha(t))} \left(\frac{\nu}{\nu_0} \right)^{\alpha(t)} \end{aligned} \quad (141)$$

where $\prime\prime$ and $\prime\prime$ indicate the signature of the Regge-trajectories. To eliminate the poles at $t < 0$ additional Γ -functions are introduced in (141). If the Pomeron trajectory is taken as $1.0 + 0.15t$ [22], negative t poles are at $\alpha = 0, -2, -4, \dots$ and therefore

$$\sin\left(\frac{\pi}{2}\alpha(t)\right) \rightarrow \sin\left(\frac{\pi}{2}\alpha(t)\right) \Gamma\left(\frac{\alpha(t)}{2}\right) \quad (142)$$

For the pion trajectory $\alpha(t) = -0.014 + 0.72t$ [22], and the negative poles are at $\alpha = -2, -4, \dots$. Regularization must be taken as

$$\sin\left(\frac{\pi}{2}\alpha(t)\right) \rightarrow \sin\left(\frac{\pi}{2}\alpha(t)\right) \Gamma\left(\frac{\alpha(t)}{2} + 1\right) \quad (143)$$

For ρ , ω and a_1 exchanges the negative poles start from $a = -1$ and therefore

$$\cos\left(\frac{\pi}{2}\alpha(t)\right) \rightarrow \cos\left(\frac{\pi}{2}\alpha(t)\right) \Gamma\left(\frac{\alpha(t)}{2} + \frac{1}{2}\right). \quad (144)$$

9.1 Single meson photoproduction due to ρ and ω exchange

In the following, the 4-vectors of the initial photon and proton are denoted as k_1 and k_2 and 4-vectors of the final state nucleon (e.g. proton) and the meson (e.g. pion) as q_1 and q_2 respectively (see Fig. 2). The photon couples to the $\pi^-\rho(770)$ system in a P-wave, and the corresponding amplitude for upper vertex is

$$A_{upper} = \varepsilon_\mu \rho_\alpha \epsilon_{\mu\alpha\beta\gamma} q_{2\beta} k_{2\gamma}, \quad (145)$$

where ρ_α is the polarization vector of ρ -meson.

Another vertex in this diagram describes the transition of




Fig. 2. The t-channel exchange diagram for single meson photoproduction

the proton and the ρ -meson into the final proton. Such a transition has the same vertex structure as the transition γ^*N to a nucleon at the lower vertex:

$$A_{lower}^{(i-)} = \bar{u}(q_1) V^{(i-)\mu}(k_1^\perp) u(k_1) \rho_\mu \quad i = 1, 2$$

$$k_{1\mu}^\perp = k_{1\nu} \left(g_{\mu\nu} - \frac{q_{1\mu} q_{1\nu}}{q_1^2} \right) = \frac{1}{2} (k_1 - k_t)_\nu \left(g_{\mu\nu} - \frac{q_{1\mu} q_{1\nu}}{q_1^2} \right) \quad (146)$$

Here $k_t = k_2 - q_2 = q_1 - k_1$ is the ρ -meson momentum of. Summing over its polarizations yields

$$\sum_{polarization} \rho_\alpha \rho_\beta = g_{\alpha\beta} - \frac{k_{t\alpha} k_{t\beta}}{k_t^2} \quad (147)$$

we obtain the following expression for the amplitude:

$$A^{i-} = \varepsilon_\mu \epsilon_{\mu\alpha\beta\gamma} q_{2\beta} k_{2\gamma} \bar{u}(q_1) V^{(i-)\alpha}(k_1^\perp) u(k_1) \quad i = 1, 2 \quad (148)$$

The same amplitude structure corresponds also to ω - exchange.

9.2 Double meson photoproduction due to ρ and ω exchange

Let us consider photoproduction of two meson (e.g. pions) due to ρ exchange in t-channel with a $1/2$ resonance in the intermediate state (see Fig. 3). In this case we should add




Fig. 3. The t-channel exchange diagram for double meson photoproduction reactions

to eq.(148) the $1/2$ propagator and the vertex for decay of this resonance into final meson and nucleon:

$$A^{i\pm} = \varepsilon_\mu \epsilon_{\mu\alpha\beta\gamma} q_{3\beta} k_{2\gamma} \bar{u}(q_1) \tilde{N}^\pm(q_{12}^\perp) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}}$$

$$V^{(i\pm)\alpha}(k_1^\perp) u(k_1), \quad i = 1, 2 \quad (149)$$

with $k_{1\mu}^\perp = k_{1\nu} (g_{\mu\nu} - (q_1 + q_2)_\mu (q_1 + q_2)_\nu / s_{12})$.

The definition of q_{12}^\perp is given in (132). The '-' amplitude corresponds to production of a $1/2^+$ intermediate state while the '+' amplitude corresponds to production of a $1/2^-$ intermediate state.

Two-meson photoproduction due to ρ exchange in t-channel with a $3/2$ resonance in the intermediate state can be easily obtained following the procedure given above. In

$$A^{i\pm} = \varepsilon_\mu \epsilon_{\mu\alpha\beta\gamma} q_{3\beta} k_{2\gamma} \bar{u}(q_1) \tilde{N}_\xi^\pm(q_{12}^\perp)$$

$$F_\chi^\xi(q_1 + q_2) V_\chi^{(i\pm)\alpha}(k_1^\perp) u(k_1) \quad i = 1, 2, 3 \quad (150)$$

the '-' amplitude corresponds to a $3/2^-$ intermediate state and '+' amplitude to $3/2^+$ intermediate state.

The examples of other t-channel and u-channel exchange amplitudes used in the analysis of the single and double meson photoproduction are given in Appendix D.

10 The cross section for photoproduction processes

The differential cross section for production of two or more particles has the form:

$$d\sigma = \frac{(2\pi)^4 |A|^2}{4\sqrt{(k_1 k_2)^2 - m_1^2 m_2^2}} d\Phi_n(k_1 + k_2, q_1, \dots, q_n) \quad (151)$$

where k_1 and k_2 are momenta of the initial particles (nucleon and γ in the case of photoproduction) and q_i are momenta of final state particles. The $d\Phi_n(k_1 + k_2, q_1, \dots, q_n)$ is the element of the n-body phase volume given by

$$d\Phi_n(k_1 + k_2, q_1, \dots, q_n) = \delta^4(k_1 + k_2 - \sum_{i=1}^n q_i) \prod_{i=1}^n \frac{d^3 q_i}{(2\pi)^3 2q_{0i}} \quad (152)$$

The photoproduction amplitude can be written as

$$A = \varepsilon_\mu \bar{u}_i A_\mu u_f \quad (153)$$

where ε_μ is the γ polarization vector and \bar{u}_i and u_f the bispinors of the initial and final state nucleon. When the γ and nucleon polarization are not measured the amplitude squared is equal to

$$|A|^2 = \frac{1}{4} \sum_{\alpha j k} A A^* = \frac{1}{4} \sum_{\alpha j k} \varepsilon_\mu^\alpha \varepsilon_\nu^\alpha \bar{u}_i^j A_\mu u_f^k \bar{u}_f^k A_\nu^{tr} u_i^j \quad (154)$$

where one averages over the polarization of the initial and sums over the polarization of the final state particles. A^{tr} is the hermitian conjugate amplitude.

For the unpolarized real photons:

$$-\sum_{\alpha} \varepsilon_\mu^\alpha \varepsilon_\nu^\alpha = g_{\mu\nu}^{\perp\perp} = g_{\mu\nu} - \frac{P_\mu P_\nu}{P^2} - \frac{k_\mu^\perp k_\nu^\perp}{k_\perp^2} \quad (155)$$

with $P = k_1 + k_2$ and

$$k_\mu^\perp = \frac{1}{2}(k_1 - k_2)_\nu g_{\mu\nu}^\perp = \frac{1}{2}(k_1 - k_2)_\nu \left(g_{\mu\nu}^\perp - \frac{P_\mu P_\nu}{P^2} \right)$$

Let us remind that in the c.m.s. with the momentum of γ being parallel to the z-axis:

$$g_{\mu\nu}^{\perp\perp} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (156)$$

The bispinors of fermions with momentum k_1 summed over polarization are convoluted (taking into account normalization (27)) and yield

$$\sum_j u^j(k_1) \bar{u}^j(k_1) = m + \hat{k}_1 \quad (157)$$

and therefore

$$|A|^2 = \frac{1}{4} g_{\mu\nu}^{\perp\perp} S p \left[(m + \hat{k}_1) A_\mu (m + \hat{k}_1) A_\nu^{tr} \right]. \quad (158)$$

In case of a polarised target the density matrix of the fermion propagator $(m + \hat{k}_1)$ must be changed to the polarization density matrix:

$$m + \hat{k}_1 \rightarrow (m + \hat{k}_1)(1 - \gamma_5 \hat{S}), \quad (159)$$

where the 4-vector S is the polarization vector of the target. When a γ is linearly polarised along the x-axis the polarization vector is: $\varepsilon_\mu = (0, 1, 0, 0)$ and we do not need to average over two polarizations. Then one has to change (158) by substituting

$$\frac{1}{2} g_{\mu\nu}^{\perp\perp} \rightarrow \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (160)$$

If one has a circular polarised beam

$$\frac{1}{2} g_{\mu\nu}^{\perp\perp} \rightarrow \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & -i & 0 \\ 0 & i & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (161)$$

11 Conclusion

In the present paper the operator expansion approach has been developed for the construction of amplitudes for pion- and photon-induced reactions. The method is relativistically invariant and can be easily applied to the construction of amplitudes with multi-body final states. For production of pseudoscalar mesons the identity of our amplitudes to the well known CGLN amplitudes is explicitly shown. The formulas are given explicitly in the form used by the Crystal Barrel at ELSA collaboration in the analysis of single and double meson photoproduction.

Acknowledgments

A. V. Anisovich and A. V. Sarantsev would like to thank the Alexander von Humboldt foundation for generous support, A.A. for a AvH fellowship and A.S. for the Friedrich-Wilhelm Bessel award. U. Thoma gratefully acknowledges a Emmy Noether grant by the Deutsche Forschungsgemeinschaft.

Appendices

A Properties of Legendre polynomials

The recurrent expression for Legendre polynomials is given by

$$P_L(z) = \frac{2L-1}{L} z P_{L-1}(z) - \frac{L-1}{L} P_{L-2}(z) \quad (162)$$

The first and second derivative of the Legendre polynomials can be expressed as

$$P'_L(z) = L \frac{P_{L-1}(z) - z P_L(z)}{1 - z^2} = (L+1) \frac{z P_L - P_{L+1}}{1 - z^2} \quad (163)$$

$$\begin{aligned} P''_L(z) &= \frac{2z P'_L(z) - L(L+1) P_L(z)}{1 - z^2} \\ &= \frac{2P'_{L+1}(z) - (L+1)(L+2) P_L(z)}{1 - z^2} \end{aligned} \quad (164)$$

Some other useful expressions given here for convenience are:

$$\begin{aligned} P'_{L-1} &= P'_L z - L P_L, \quad P'_{L+1} = P'_L z + (L+1) P_L \\ P'_{L+1} - P'_{L-1} &= (2L+1) P_L, \\ P''_{L+1} - P''_{L-1} &= (2L+1) P'_L \end{aligned}$$

B Properties of angular momentum operators

In the following we list useful properties of angular momentum operators.

$$\begin{aligned} X_{\mu\alpha_1\dots\alpha_n}^{(n+1)}(q_\perp) X_{\alpha_1\dots\alpha_n}^{(n)}(k_\perp) &= \\ \frac{\alpha_n}{n+1} (\sqrt{k_\perp^2})^n (\sqrt{q_\perp^2})^{n+1} &\left[-\frac{k_{1\mu}}{\sqrt{k_\perp^2}} P'_n + \frac{q_{1\mu}}{\sqrt{q_\perp^2}} P'_{n+1} \right] \end{aligned} \quad (165)$$

$$\begin{aligned} X_{\mu\alpha_2\dots\alpha_n}^{(n)}(q_\perp) X_{\nu\alpha_2\dots\alpha_n}^{(n)}(k_\perp) &= \\ \frac{\alpha_{n-1}}{n^2} (\sqrt{k_\perp^2})^n (\sqrt{q_\perp^2})^n &\left[g_{\mu\nu}^{\perp} P'_{n-1} - \left(\frac{q_\mu^\perp q_\nu^\perp}{q_\perp^2} + \frac{k_\mu^\perp k_\nu^\perp}{k_\perp^2} \right) P'_n + \right. \\ \frac{1}{2} \left(\frac{q_\mu^\perp k_\nu^\perp + k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2} \sqrt{q_\perp^2}} \right) &(P'_n + 2z P''_n) + \\ \left. \frac{2n-1}{2} \left(\frac{q_\mu^\perp k_\nu^\perp - k_\mu^\perp q_\nu^\perp}{\sqrt{k_\perp^2} \sqrt{q_\perp^2}} \right) P'_n \right] \end{aligned} \quad (166)$$

$$\begin{aligned} X_{\mu\alpha_1\dots\alpha_n}^{(n+2)}(q_\perp) X_{\alpha_1\dots\alpha_n}^{(n)}(k_\perp) &= \\ \frac{2}{3} \frac{\alpha_n}{(n+1)(n+2)} (\sqrt{k_\perp^2})^n (\sqrt{q_\perp^2})^{n+2} &\left(\frac{X_{\mu\nu}^{(2)}(q_\perp)}{q_\perp^2} P''_{n+2} + \right. \\ \left. \frac{X_{\mu\nu}^{(2)}(k_\perp)}{k_\perp^2} P''_n - \frac{3}{2} \frac{k_\mu^\perp q_\nu^\perp + k_\nu^\perp q_\mu^\perp - \frac{2}{3} g_{\mu\nu}^{\perp} (k_\perp^\perp q_\perp^\perp)}{\sqrt{k_\perp^2} \sqrt{q_\perp^2}} P''_{n+1} \right) \end{aligned} \quad (167)$$

$$\begin{aligned} X_{\alpha\gamma_2\dots\gamma_n}^{(n)}(q_\perp) O^{\tau\gamma_2\dots\gamma_n} X_{\xi\beta_2\dots\beta_n}^{(n)}(k_\perp) &= \\ X_{\alpha\gamma_2\dots\gamma_n}^{(n)}(q_\perp) \frac{g_{\tau\mu}}{n} X_{\xi\gamma_2\dots\gamma_n}^{(n)}(k_\perp) &+ \\ + \frac{n-1}{n} X_{\alpha\mu\gamma_3\dots\gamma_n}^n(q_\perp) X_{\xi\tau\gamma_3\dots\gamma_n}^{(n)}(k_\perp) &- \\ \frac{2(n-1)}{n(2n-1)} X_{\alpha\tau\gamma_3\dots\gamma_n}^{(n)}(q_\perp) X_{\xi\mu\gamma_3\dots\gamma_n}^{(n)}(k_\perp) \end{aligned} \quad (168)$$

C Blatt-Weisskopf formfactors

If a resonance with radius r decays into two particles with (squared) momentum k^2 :

$$k^2 = \frac{(s - (m_1 + m_2)^2)(s - (m_1 - m_2)^2)}{4s}, \quad (169)$$

where s is total energy and m_1 and m_2 are masses of the final particles, then the first few expressions for formfactors $F(L, k^2, r)$ are

$$\begin{aligned} F(0, k^2, r) &= 1 \\ F(1, k^2, r) &= \frac{\sqrt{(x+1)}}{r} \\ F(2, k^2, r) &= \frac{\sqrt{(x^2 + 3x + 9)}}{r^2} \\ F(3, k^2, r) &= \frac{\sqrt{(x^3 + 6x^2 + 45x + 225)}}{r^3} \\ F(4, k^2, r) &= \frac{\sqrt{(x^4 + 10x^3 + 135x^2 + 1575x + 11025)}}{r^4}, \end{aligned} \quad (170)$$

where $x = k^2 r^2$. Remember that $r(\text{GeV}^{-1}) = r(fm)/(0.1973(fm\text{GeV}))$.

D Structure of amplitudes for t-channel and u-channel exchanges

D1 t-channel amplitudes

For the photoproduction of a single neutral pion, ρ and ω exchanges play a significant role. The exchange of a π^0 is forbidden since the photon does not couple to a neutral pion. When charged pions are produced the pion exchange diagram can play an important role. The upper vertex function for pion exchange is

$$\begin{aligned} A_{upper} &= \varepsilon_\mu \frac{1}{2} (k_2 + k_t)_\nu \left(g_{\mu\nu} - \frac{q_{2\mu} q_{2\nu}}{m_\pi^2} \right) = \\ \varepsilon_\mu k_{t\nu} \left(g_{\mu\nu} - \frac{q_{2\mu} q_{2\nu}}{m_\pi^2} \right) \end{aligned} \quad (171)$$

The lower vertex function is described by a $N \rightarrow \pi N$ transition. Thus

$$A = \varepsilon_\mu k_{t\nu} \left(g_{\mu\nu} - \frac{q_{2\mu} q_{2\nu}}{m_\pi^2} \right) \bar{u}(q_1) N^-(k_1^\perp) u(k_1) \quad (172)$$

Remember that for single meson production

$$k_{1\mu}^\perp = \frac{1}{2} (k_1 - k_t)_\nu \left(g_{\mu\nu} - \frac{q_{1\mu} q_{1\nu}}{q_1^2} \right) = k_{1\nu} \left(g_{\mu\nu} - \frac{q_{1\mu} q_{1\nu}}{q_1^2} \right) \quad (173)$$

This expression can be easily extended to the case of double meson photoproduction. If the intermediate baryon has spin 1/2 one obtains

$$\begin{aligned} A^\pm &= \varepsilon_\mu k_{t\nu} \left(g_{\mu\nu} - \frac{q_{2\mu} q_{2\nu}}{m_\pi^2} \right) \\ \bar{u}(q_1) \tilde{N}^\pm(q_{12}^\perp) \frac{\hat{q}_1 + \hat{q}_2 + \sqrt{s_{12}}}{2\sqrt{s_{12}}} N^\pm(k_1^\perp) u(k_1) \end{aligned} \quad (174)$$

Here the '-' amplitude corresponds to a $1/2^+$ intermediate state, the '+' to a $1/2^-$ state,

$$k_{1\mu}^\perp = k_{1\nu} (g_{\mu\nu} - (q_1 + q_2)_\mu (q_1 + q_2)_\nu / s_{12}), \quad (175)$$

the definition of q_{12}^\perp is given in eqn.(132) and the notation of momenta is shown in Fig. 3.

For an intermediate resonance with spin $J = L \pm 1/2$ the amplitude structure reads

$$A^\pm = \varepsilon_\mu k_{t\nu} \left(g_{\mu\nu} - \frac{q_{2\mu} q_{2\nu}}{m_\pi^2} \right) \bar{u}(q_1) \tilde{N}_{\alpha_1 \alpha_2 \dots \alpha_n}^\pm(q_{12}^\perp) F_{\beta_1 \beta_2 \dots \beta_L}^{\alpha_1 \alpha_2 \dots \alpha_n}(q_1 + q_2) N_{\beta_1 \beta_2 \dots \beta_L}^\pm(k_1^\perp) u(k_1) \quad (176)$$

The upper vertex for ρ -meson production due to pion exchange has the following structure

$$A_{upper} = \varepsilon_\mu \varepsilon_{\mu\alpha\beta\gamma} \frac{1}{2} (q_3 - q_2)_\alpha q_{2\beta} k_{2\gamma} = \varepsilon_\mu \varepsilon_{\mu\alpha\beta\gamma} q_{3\alpha} q_{2\beta} k_{2\gamma}, \quad (177)$$

while the lower vertex has the same structure as the πN scattering amplitude. Therefore

$$A = \varepsilon_\mu \varepsilon_{\mu\alpha\beta\gamma} q_{3\alpha} q_{2\beta} k_{2\gamma} \bar{u}(q_1) N^-(k_1^\perp) u(k_1) \quad (178)$$

Here k_1^\perp is given by eq.(173).

The ρ meson can also be produced by Pomeron or f_0 exchange. The upper vertex for such a case is $\varepsilon_\mu \frac{1}{2} (q_3 - q_2)_\mu$ and the amplitude is equal to

$$A = \varepsilon_\mu \frac{1}{2} (q_3 - q_2)_\mu \bar{u}(q_1) N^+(k_1^\perp) u(k_1) \quad (179)$$

The next amplitude which we consider is the f_0 production due to ρ (or ω) t-channel exchange. Such an amplitude has the structure:

$$A^{i-} = \varepsilon_\mu \left(g_{\mu\nu} - \frac{k_{t\mu} k_{t\nu}}{k_t^2} \right) \bar{u}(q_1) V_\nu^{(i-)}(k_1^\perp) u(k_1) \quad i = 1, 2 \quad (180)$$

D2 u-channel amplitudes




Fig. 4. The u-channel exchange diagram for photoproduction of single mesons

Apart from meson exchange amplitudes (which we define as t-channel exchanges), mesons can be produced from baryon exchange in the u-channel. An example of such a diagram is given in Fig. 4. For nucleon exchange, the vertex for meson production (the lower vertex) is defined by

$$\bar{u}(k_u) N^-(q_2^\perp) u(k_1) \quad k_u = k_1 - q_2. \quad (181)$$

Here the N^- vertex describes the production of a pseudoscalar meson. Further,

$$q_{2\mu}^\perp = \left(g_{\mu\nu} - \frac{k_{1\mu} k_{1\nu}}{m_p^2} \right) \frac{1}{2} (q_2 - k_u)_\nu = \left(g_{\mu\nu} - \frac{k_{1\mu} k_{1\nu}}{m_p^2} \right) q_{2\nu}. \quad (182)$$




Fig. 5. A u-channel exchange diagram for production of a baryon resonance in photoproduction of two mesons

If the reaction is induced by a meson the upper vertex has the same structure

$$\bar{u}(q_1) N^-(k_2^\perp) u(k_u) \quad (183)$$

where

$$k_{2\mu}^\perp = \left(g_{\mu\nu} - \frac{q_{1\mu} q_{1\nu}}{m_p^2} \right) \frac{1}{2} (k_2 - k_u)_\nu = \left(g_{\mu\nu} - \frac{q_{1\mu} q_{1\nu}}{m_p^2} \right) k_{2\nu}$$

The angular dependent part of the amplitude for the nucleon exchange diagram is

$$A = \bar{u}(q_1) N^-(k_2^\perp) \frac{m_p + \hat{k}_u}{m_p^2 - k_u^2} N^-(q_2^\perp) u(k_1) \quad (184)$$

In the case of photoproduction the upper vertex is defined by $V_\mu^{(i-)}$:

$$A^i = \varepsilon_\mu \bar{u}(q_1) V_\mu^{(i-)}(k_2^\perp) \frac{m_p + \hat{k}_u}{m_p^2 - k_u^2} N^-(q_2^\perp) u(k_1), \quad i = 1, 2 \quad (185)$$

The production of 0^{++} states in double meson production can be obtained from eqs.(184,185) by replacing $N^-(q_2^\perp)$ by $N^+(q_2^\perp)$.

In the case when a baryon resonance with $J = L \pm 1/2$ is produced in the intermediate state (see Fig. 5) the amplitude for meson induced reaction has the structure

$$A = \bar{u}(q_1) N_{\alpha_1 \alpha_2 \dots \alpha_L}^{\pm} (q_{12}^{\perp}) F_{\beta_1 \beta_2 \dots \beta_L}^{\alpha_1 \alpha_2 \dots \alpha_L} (q_1 + q_2) N_{\beta_1 \beta_2 \dots \beta_L}^{\pm} (k_2^{\perp}) \frac{m_p + \hat{k}_u}{m_p^2 - k_u^2} N^{-}(q_3^{\perp}) u(k_1) \quad (186)$$

and for γ^* induced reactions

$$A^{i\pm} = \bar{u}(q_1) V_{\alpha_1 \alpha_2 \dots \alpha_L}^{(i\pm)} (q_{12}^{\perp}) F_{\beta_1 \beta_2 \dots \beta_L}^{\alpha_1 \alpha_2 \dots \alpha_L} (q_1 + q_2) N_{\beta_1 \beta_2 \dots \beta_L}^{\pm} (k_2^{\perp}) \frac{m_p + \hat{k}_u}{m_p^2 - k_u^2} N^{-}(q_3^{\perp}) u(k_1) \quad (187)$$

References

1. C. Amsler and F. E. Close, Phys. Lett. B **353** (1995) 385.
2. V. V. Anisovich and A. V. Sarantsev, Nucl. Phys. Proc. Suppl. **54A** (1997) 367.
3. P. Minkowski and W. Ochs, Eur. Phys. J. C **9** (1999) 283.
4. E. Klempt, *Meson spectroscopy: glueballs, hybrids, and q anti- q mesons*, hep-ex/0101031 (2001).
5. A. V. Anisovich, V. V. Anisovich and A. V. Sarantsev, Phys. Rev. D **62**, 051502 (2000) [arXiv:hep-ph/0003113].
6. E. Klempt, Phys. Rev. C **66** (2002) 058201.
7. V. V. Anisovich, D. V. Bugg, A. V. Sarantsev and B. S. Zou, Phys. Atom. Nucl. **57** (1994) 1595 [Yad. Fiz. **57** (1994) 1666].
8. V.V. Anisovich and L.G. Dakhno, JETP **44** (1963) 198; V.V. Anisovich and A.A. Anselm, UFN **88**, 287 (1965), [Sov. Phys. Uspekhi **88**, 177 (1966)].
9. C. Zemach, Phys. Rev. **97** (1965) B97; **97** (1965) B109.
10. A.V. Anisovich and A.V. Sarantsev, Sov. J. Nucl. Phys. **55** (1992) 1200.
11. V. V. Anisovich, M. N. Kobrinsky, D. I. Melikhov and A. V. Sarantsev, Nucl. Phys. A **544** (1992) 747.
12. A.V. Anisovich and V.A. Sadovnikova, Sov. J. Nucl. Phys. **55** (1992) 1483; Eur. Phys. J. **A2** (1998) 199.
13. S.U. Chung, Phys. Rev. **D57** (1998) 431.
14. A. V. Anisovich, V. V. Anisovich, V. N. Markov, M. A. Matveev and A. V. Sarantsev, J. Phys. G **28** (2002) 15.
15. A. V. Anisovich, V. V. Anisovich, M. A. Matveev and V. A. Nikonov, Phys. Atom. Nucl. **66** (2003) 914 [Yad. Fiz. **66** (2003) 946].
16. B. Krusche *et al.*, Phys. Rev. Lett. **74** (1995) 3736. R. Beck *et al.*, Phys. Rev. Lett. **78** (1997) 606.
17. J. Ajaka *et al.*, Phys. Rev. Lett. **81** (1998) 1797. D. Rebreyend *et al.*, Nucl. Phys. A **663** (2000) 436. F. Renard *et al.*, Phys. Lett. B **528** (2002) 215. D. Rebreyend, private communication, 2004.
18. K. H. Glander *et al.*, Eur. Phys. J. A **19** (2004) 251. J. Barth *et al.*, Eur. Phys. J. A. **18** (2003) 117. J. Barth *et al.*, Phys. Lett. B **572** (2003) 127. J. Barth *et al.*, Eur. Phys. J. A **17** (2003) 269. S. Goers *et al.*, Phys. Lett. B **464** (1999) 331. R. Plötzke *et al.*, Phys. Lett. B **444** (1998) 555. C. Bennhold *et al.*, Nucl. Phys. A **639** (1998) 209.
19. O. Bartholomy *et al.*, submitted to PRL. V. Crede *et al.*, submitted to PRL.
20. V. Kubarovsky *et al.*, Phys. Rev. Lett. **92** (2004) 032001 [Erratum-ibid. **92** (2004) 049902]. J. W. C. McNabb *et al.*, Phys. Rev. C **69** (2004) 042201. S. Stepanyan *et al.*, Phys. Rev. Lett. **91** (2003) 252001. M. Ripani *et al.*, Phys. Rev. Lett. **91** (2003) 022002. M. Dugger *et al.*, Phys. Rev. Lett. **89** (2002) 222002 [Erratum-ibid. **89** (2002) 249904]. R. Thompson *et al.*, Phys. Rev. Lett. **86** (2001) 1702. S. P. Barrow *et al.*, Phys. Rev. C **64** (2001) 044601.
21. D. Drechsel, O. Hanstein, S. S. Kamalov and L. Tiator, Nucl. Phys. A **645** (1999) 145.
22. A.B. Kaidalov and B.M. Karnakov, Yad. Fiz. **11** (1970) 216. G.D. Alkhozov, V.V. Anisovich and P.E. Volkovitsky, "Diffractive interaction of high energy hadrons on nuclei", Chapter I, "Science", Leningrad, 1991.