

# Covariant tensor formalism for partial wave analyses of $\psi$ decay to mesons

**B.S. Zou<sup>a,b,1</sup> and D.V.Bugg<sup>c</sup>**

a) CCAST (World Laboratory), P.O. Box 8730, Beijing 100080

b) Institute of High Energy Physics, CAS, P.O. Box 918(4), Beijing 100039, China

c) Queen Mary College, London, UK

## Abstract

$J/\psi$  and  $\psi'$  decay to mesons are a good place to look for glueballs, hybrids and for extracting strange and nonstrange components in mesons. Abundant  $J/\psi$  and  $\psi'$  events have been collected at the Beijing Electron Positron Collider (BEPC). More data will be collected at upgraded BEPC and CLEO-C. Here we provide explicit PWA formulae for many interesting channels in the covariant tensor formalism.

## 1 Introduction

High statistics data have appeared from BES for  $J/\psi$  decays and will soon be available also for  $\psi'$  decays. Further high statistics data are expected from CLEO[1]. It is convenient to have a uniform approach to partial wave analyses. Here we provide one such approach using covariant tensor formalism. A similar approach has been used in analyzing other reactions[2, 3, 4]. We provide formulae documenting those which have been used for a number of channels already published by BES[5, 6, 7, 8, 9] and extend them to further channels being prepared for publication. This list of reactions is not exhaustive, but formulae are readily extended to other cases following the same methods.

Reactions fall into two categories: non-radiative decays, where final-state particles are pions or kaons; all polarization information is then available in the form of angular distributions. Reactions of this type are discussed in Section 2. This formalism extends also to final states containing the  $\omega$ , where polarization information is measured fully by the decay  $\omega \rightarrow \pi^+\pi^-\pi^0$ . The second class of reactions consists of radiative decays, e.g.  $J/\psi \rightarrow \gamma\pi^+\pi^-$ . For this class, differential cross sections need to be summed over the unmeasured helicities of the photon, incorporating the knowledge that the photon is transverse. These reactions are considered in Section 3.

---

<sup>1</sup>zoubs@mail.ihep.ac.cn

## 2 Formalism for $\psi$ non-radiative decay to mesons

The general form for the decay amplitude of a vector meson  $\psi$  with spin projection of  $m$  is

$$A = \psi_\mu(m)A^\mu = \psi_\mu(m) \sum_i \Lambda_i U_i^\mu \quad (1)$$

where  $\psi_\mu(m)$  is the polarization vector of the  $\psi$ ;  $U_i^\mu$  is the  $i$ -th partial wave amplitude with coupling strength determined by a complex parameter  $\Lambda_i$ . The polarization vector satisfies

$$\sum_{m=1}^3 \psi^\mu(m)\psi^{*\nu}(m) = -g^{\mu\nu} + \frac{p_\psi^\mu p_\psi^\nu}{p_\psi^2} \equiv -\tilde{g}^{\mu\nu}(p_\psi). \quad (2)$$

For  $\psi$  production from  $e^+e^-$  annihilation, the electrons are highly relativistic, with the result that  $J_z = \pm 1$ . If we take the beam direction to be the  $z$ -axis, this limits  $m$  to 1 and 2, i.e. components along  $x$  and  $y$ . Then the differential cross section for decay to an  $n$ -body final state is:

$$\frac{d\sigma}{d\Phi_n} = \frac{(2\pi)^4}{2M_\psi} \cdot \frac{1}{2} \sum_{m=1}^2 \psi_\mu(m)A^\mu \psi_{\mu'}^*(m)A^{*\mu'} \quad (3)$$

where  $M_\psi$  is the mass of  $\psi$  and  $d\Phi_n$  is the standard element of  $n$ -body phase space given by

$$d\Phi_n(p_\psi; p_1, \dots, p_n) = \delta^4(p_\psi - \sum_{i=1}^n p_i) \prod_{i=1}^n \frac{d^3\mathbf{p}_i}{(2\pi)^3 2E_i}. \quad (4)$$

Note that

$$\sum_{m=1}^2 \psi_\mu(m)\psi_{\mu'}^*(m) = \delta_{\mu\mu'}(\delta_{\mu 1} + \delta_{\mu 2}), \quad (5)$$

so we have

$$\frac{d\sigma}{d\Phi_n} = \frac{1}{2} \sum_{\mu=1}^2 A^\mu A^{*\mu} = \frac{1}{2} \sum_{i,j} \Lambda_i \Lambda_j^* \sum_{\mu=1}^2 U_i^\mu U_j^{*\mu} \equiv \sum_{i,j} P_{ij} \cdot F_{ij} \quad (6)$$

where

$$P_{ij} = P_{ji}^* = \Lambda_i \Lambda_j^*, \quad (7)$$

$$F_{ij} = F_{ji}^* = \frac{1}{2} \sum_{\mu=1}^2 U_i^\mu U_j^{*\mu}. \quad (8)$$

We construct the partial wave amplitudes  $U_i^\mu$  in the covariant Rarita-Schwinger tensor formalism [10]. As in Ref. [11], we use pure orbital angular momentum covariant tensors  $\tilde{t}_{\mu_1 \dots \mu_l}^{(l)}$  and covariant spin wave functions  $\phi_{\mu_1 \dots \mu_S}$  together with operators  $g_{\mu\nu}$ ,  $\epsilon_{\mu\nu\lambda\sigma}$  and

momenta of parent particles. For a process  $a \rightarrow bc$ , the covariant tensors  $\tilde{t}_{\mu_1 \dots \mu_l}^{(l)}$  for final states of pure orbital angular momentum  $l$  are constructed from relevant momenta  $p_a$ ,  $p_b$  and  $p_c$  [11]

$$\tilde{t}^{(0)} = 1, \quad (9)$$

$$\tilde{t}_\mu^{(1)} = \tilde{g}_{\mu\nu}(p_a)r^\nu B_1(Q_{abc}) \equiv \tilde{r}_\mu B_1(Q_{abc}), \quad (10)$$

$$\tilde{t}_{\mu\nu}^{(2)} = [\tilde{r}_\mu \tilde{r}_\nu - \frac{1}{3}(\tilde{r} \cdot \tilde{r})\tilde{g}_{\mu\nu}(p_a)] B_2(Q_{abc}), \quad (11)$$

$$\tilde{t}_{\mu\nu\lambda}^{(3)} = [\tilde{r}_\mu \tilde{r}_\nu \tilde{r}_\lambda - \frac{1}{5}(\tilde{r} \cdot \tilde{r})(\tilde{g}_{\mu\nu}(p_a)\tilde{r}_\lambda + \tilde{g}_{\nu\lambda}(p_a)\tilde{r}_\mu + \tilde{g}_{\lambda\mu}(p_a)\tilde{r}_\nu)] B_3(Q_{abc}), \quad (12)$$

...

with  $r = p_b - p_c$ . The term  $(\tilde{r} \cdot \tilde{r})$  is the dot-product of 4-vectors:  $\tilde{r}_0\tilde{r}_0 - \tilde{r}_1\tilde{r}_1 - \tilde{r}_2\tilde{r}_2 - \tilde{r}_3\tilde{r}_3$ , and makes  $\tilde{t}_{\mu\nu}^{(2)}$  traceless. Likewise  $\tilde{t}^{(3)}$  is constructed to be traceless.  $Q_{abc}$  is the magnitude of  $\mathbf{p}_b$  or  $\mathbf{p}_c$  in the rest system of  $a$ , where

$$Q_{abc}^2 = \frac{(s_a + s_b - s_c)^2}{4s_a} - s_b \quad (13)$$

with  $s_a = E_a^2 - \mathbf{p}_a^2$ . Then  $\tilde{t}_{\mu_1 \dots \mu_l}^{(l)}$  contains the angular distribution function multiplied by a Blatt-Weisskopf barrier factor [11, 12]  $Q_{abc}^l B_l(Q_{abc})$ . Explicitly

$$B_1(Q_{abc}) = \sqrt{\frac{2}{Q_{abc}^2 + Q_0^2}}, \quad (14)$$

$$B_2(Q_{abc}) = \sqrt{\frac{13}{Q_{abc}^4 + 3Q_{abc}^2 Q_0^2 + 9Q_0^4}}, \quad (15)$$

$$B_3(Q_{abc}) = \sqrt{\frac{277}{Q_{abc}^6 + 6Q_{abc}^4 Q_0^2 + 45Q_{abc}^2 Q_0^4 + 225Q_0^6}}, \quad (16)$$

$$B_4(Q_{abc}) = \sqrt{\frac{12746}{Q_{abc}^8 + 10Q_{abc}^6 Q_0^2 + 135Q_{abc}^4 Q_0^4 + 1575Q_{abc}^2 Q_0^6 + 11025Q_0^8}} \quad (17)$$

Here  $Q_0$  is a hadron ‘‘scale’’ parameter  $Q_0 = 0.197321/R$  GeV/ $c$ , where  $R$  is the radius of the centrifugal barrier in fm. We remark that in these Blatt-Weisskopf factors, the approximation is made that the centrifugal barrier may be replaced by a square well of radius  $R$ .

If  $a$  is an intermediate resonance decaying into  $bc$ , one needs to introduce into the amplitude a Breit-Wigner propagator denoted by  $f_{(bc)}^{(a)}$ :

$$f_{(bc)}^{(a)} = \frac{1}{m_a^2 - s_{bc} - im_a \Gamma_a}; \quad (18)$$

here  $s_{bc} = (p_b + p_c)^2$  is the invariant mass-squared of  $b$  and  $c$ ;  $m_a, \Gamma_a$  are the resonance mass and width.

We outline now some further general features of notation, taking as an example the two-step process  $J/\psi \rightarrow \rho_{12}\pi_3, \rho_{12} \rightarrow \pi_1\pi_2$ . In the first step we denote the orbital angular momentum by  $L$ ; in this example  $L = 1$ . In the second step, we denote the orbital angular momentum by  $\ell$ , which is again 1 in this case. The tensor describing the first step will be denoted by  $\tilde{T}_{\mu_1 \dots \mu_L}^{(L)}$ . The tensor describing the second step will be denoted by  $\tilde{t}_{\mu_1 \dots \mu_\ell}^{(\ell)}$ . The orbital angular momentum is constructed in terms of relative momenta, so it is convenient to define  $q_{(ij)} = p_i - p_j$ .

Some expressions depend also on the total momentum of the  $ij$  pair:  $p_{(ij)} = p_i + p_j$ . When one wants to combine two angular momenta ( $\mathbf{j}_b$  and  $\mathbf{j}_c$ ) into a total angular momentum  $\mathbf{j}_a$ , if  $j_a + j_b + j_c$  is an odd number, then a combination  $\epsilon_{\mu\nu\lambda\sigma} p_a^\mu$  with  $p_a$  the momentum of the parent particle is needed; otherwise it is not needed.

Projection operators will be a useful general tool in constructing expressions. For a meson  $a$  with spin  $S$  and corresponding spin wave function  $\phi_{\mu_1 \dots \mu_S}(p_a, m)$ , what we usually need to use in constructing amplitudes is its spin projection operator  $P_{\mu_1 \dots \mu_S \mu'_1 \dots \mu'_S}^{(S)}(p_a)$ .

$$P_{\mu\mu'}^{(1)}(p_a) = \sum_m \phi_\mu(p_a, m) \phi_{\mu'}^*(p_a, m) = -g_{\mu\mu'} + \frac{p_{a\mu} p_{a\mu'}}{p_a^2} \equiv -\tilde{g}_{\mu\mu'}(p_a), \quad (19)$$

$$P_{\mu\nu\mu'\nu'}^{(2)}(p_a) = \sum_m \phi_{\mu\nu}(p_a, m) \phi_{\mu'\nu'}^*(p_a, m) = \frac{1}{2}(\tilde{g}_{\mu\mu'}\tilde{g}_{\nu\nu'} + \tilde{g}_{\mu\nu'}\tilde{g}_{\nu\mu'}) - \frac{1}{3}\tilde{g}_{\mu\nu}\tilde{g}_{\mu'\nu'}, \quad (20)$$

$$\begin{aligned} P_{\mu\nu\lambda\mu'\nu'\lambda'}^{(3)}(p_a) &= \sum_m \phi_{\mu\nu\lambda}(p_a, m) \phi_{\mu'\nu'\lambda'}^*(p_a, m) \\ &= -\frac{1}{6}(\tilde{g}_{\mu\mu'}\tilde{g}_{\nu\nu'}\tilde{g}_{\lambda\lambda'} + \tilde{g}_{\mu\mu'}\tilde{g}_{\nu\lambda'}\tilde{g}_{\lambda\nu'} + \tilde{g}_{\mu\nu'}\tilde{g}_{\nu\mu'}\tilde{g}_{\lambda\lambda'} \\ &\quad + \tilde{g}_{\mu\nu'}\tilde{g}_{\nu\lambda'}\tilde{g}_{\lambda\mu'} + \tilde{g}_{\mu\lambda'}\tilde{g}_{\nu\nu'}\tilde{g}_{\lambda\mu'} + \tilde{g}_{\mu\lambda'}\tilde{g}_{\nu\mu'}\tilde{g}_{\lambda\nu'}) \\ &\quad + \frac{1}{15}(\tilde{g}_{\mu\nu}\tilde{g}_{\mu'\nu'}\tilde{g}_{\lambda\lambda'} + \tilde{g}_{\mu\nu}\tilde{g}_{\nu'\lambda'}\tilde{g}_{\lambda\mu'} + \tilde{g}_{\mu\nu}\tilde{g}_{\mu'\lambda'}\tilde{g}_{\lambda\nu'} \\ &\quad + \tilde{g}_{\mu\lambda}\tilde{g}_{\mu'\lambda'}\tilde{g}_{\nu\nu'} + \tilde{g}_{\mu\lambda}\tilde{g}_{\mu'\nu'}\tilde{g}_{\nu\lambda'} + \tilde{g}_{\mu\lambda}\tilde{g}_{\nu'\lambda'}\tilde{g}_{\nu\mu'} \\ &\quad + \tilde{g}_{\nu\lambda}\tilde{g}_{\nu'\lambda'}\tilde{g}_{\mu\mu'} + \tilde{g}_{\nu\lambda}\tilde{g}_{\mu'\nu'}\tilde{g}_{\mu\lambda'} + \tilde{g}_{\nu\lambda}\tilde{g}_{\mu'\lambda'}\tilde{g}_{\mu\nu'}), \end{aligned} \quad (21)$$

$$\begin{aligned} P_{\mu\nu\lambda\sigma\mu'\nu'\lambda'\sigma'}^{(4)}(p_a) &= \sum_m \phi_{\mu\nu\lambda\sigma}(p_a, m) \phi_{\mu'\nu'\lambda'\sigma'}^*(p_a, m) \\ &= \frac{1}{24}[\tilde{g}_{\mu\mu'}\tilde{g}_{\nu\nu'}\tilde{g}_{\lambda\lambda'}\tilde{g}_{\sigma\sigma'} + \dots (\mu', \nu', \lambda', \sigma' \text{ permutation, 24 terms})] \\ &\quad - \frac{1}{84}[\tilde{g}_{\mu\nu}\tilde{g}_{\mu'\nu'}\tilde{g}_{\lambda\lambda'}\tilde{g}_{\sigma\sigma'} + \dots (\mu, \nu, \lambda, \sigma \text{ permutation,} \\ &\quad \mu', \nu', \lambda', \sigma' \text{ permutation, 72 terms})] \\ &\quad + \frac{1}{105}(\tilde{g}_{\mu\nu}\tilde{g}_{\lambda\sigma} + \tilde{g}_{\mu\lambda}\tilde{g}_{\nu\sigma} + \tilde{g}_{\mu\sigma}\tilde{g}_{\nu\lambda})(\tilde{g}_{\mu'\nu'}\tilde{g}_{\lambda'\sigma'} + \tilde{g}_{\mu'\lambda'}\tilde{g}_{\nu'\sigma'} + \tilde{g}_{\mu'\sigma'}\tilde{g}_{\nu'\lambda'}). \end{aligned}$$

(22)

Note that

$$\tilde{t}_{\mu_1 \dots \mu_L}^{(L)} = (-1)^L P_{\mu_1 \dots \mu_L \mu'_1 \dots \mu'_L}^{(L)} r^{\mu'_1} \dots r^{\mu'_L} B_L(Q_{abc}). \quad (23)$$

We come now to specific examples of reactions.

## 2.1 $\psi \rightarrow \pi^+ \pi^- \pi^0$

For three isospin 1 particles coupling to an isospin zero particle, the only possible coupling for isospin conservation is  $(\mathbf{I}_1 \times \mathbf{I}_2) \cdot \mathbf{I}_3$ , which is fully anti-symmetric in particles 1,2,3. This demands that the angular dependent part should also be fully anti-symmetric for 1,2,3, in order to make the overall amplitude symmetric. For  $\psi \rightarrow \pi^+ \pi^- \pi^0$ , any two pions are limited to an overall isospin 1 and hence can only be negative parity states with  $J$  odd, *i.e.*,  $J^P = 1^-, 3^-, 5^-$  *etc.*

For  $\psi \rightarrow \rho(1^-) \pi \rightarrow \pi^+ \pi^- \pi^0$ ,  $\psi$  decays to  $\rho\pi$  in a P-wave; then  $\rho$  decays to  $\pi\pi$  also in P-wave, hence the amplitude for the two step process is

$$\begin{aligned} U_\rho^\mu &= (\mathbf{I}_1 \times \mathbf{I}_2) \cdot \mathbf{I}_3 \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma \tilde{T}_{(\rho 3)}^{(1)\nu} \tilde{t}_{(12)}^{(1)\lambda} f_{(12)}^{(\rho)} + (1 \leftrightarrow 3) + (2 \leftrightarrow 3) \\ &= 4i \epsilon_{\mu\nu\lambda\sigma} p_1^\nu p_2^\lambda p_3^\sigma \left[ B_1(Q_{\psi\rho 3}) f_{(12)}^{(\rho)} B_1(Q_{\rho 12}) + B_1(Q_{\psi\rho 2}) f_{(13)}^{(\rho)} B_1(Q_{\rho 13}) \right. \\ &\quad \left. + B_1(Q_{\psi\rho 1}) f_{(23)}^{(\rho)} B_1(Q_{\rho 23}) \right]. \end{aligned} \quad (24)$$

Here we use the convention  $\mathbf{I}_1 = (\frac{-1}{\sqrt{2}}, \frac{-i}{\sqrt{2}}, 0)$  for  $\pi^+$ ,  $\mathbf{I}_2 = (\frac{1}{\sqrt{2}}, \frac{-i}{\sqrt{2}}, 0)$  for  $\pi^-$  and  $\mathbf{I}_3 = (0, 0, 1)$  for  $\pi^0$ . This gives  $(\mathbf{I}_1 \times \mathbf{I}_2) \cdot \mathbf{I}_3 = -i$ .

The amplitude can be further simplified in the  $\psi$  rest system as

$$\begin{aligned} U_\rho^\mu &= 4i M_\psi \epsilon_{\mu\nu\lambda\sigma} p_1^\nu p_2^\lambda \left[ B_1(Q_{\psi\rho 3}) f_{(12)}^{(\rho)} B_1(Q_{\rho 12}) + B_1(Q_{\psi\rho 2}) f_{(13)}^{(\rho)} B_1(Q_{\rho 13}) \right. \\ &\quad \left. + B_1(Q_{\psi\rho 1}) f_{(23)}^{(\rho)} B_1(Q_{\rho 23}) \right]. \end{aligned} \quad (25)$$

For any other  $1^-$  intermediate state  $\rho'$ , one can get the corresponding amplitude by simply replacing the Breit-Wigner component  $f^{(\rho)}$  by  $f^{(\rho')}$ .

For  $\psi \rightarrow \rho_3(3^-) \pi \rightarrow \pi^+ \pi^- \pi^0$ ,  $\psi$  decays to  $\rho_3\pi$  in F-wave; then  $\rho_3$  decays to  $\pi\pi$  also in F-wave; the amplitude is

$$\begin{aligned} U_{\rho_3}^\mu &= (\mathbf{I}_1 \times \mathbf{I}_2) \cdot \mathbf{I}_3 \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma \tilde{T}_{(\rho_3 3)}^{(3)\nu\alpha\beta} \tilde{t}_{(12)\alpha\beta}^{(3)\lambda} \cdot f_{(12)}^{(\rho_3)} + (1 \leftrightarrow 3) + (2 \leftrightarrow 3) \\ &= -i \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma [\tilde{T}_{(\rho_3 3)}^{(3)\nu\alpha\beta} \tilde{t}_{(12)\alpha\beta}^{(3)\lambda} \cdot f_{(12)}^{(\rho_3)} - (1 \leftrightarrow 3) - (2 \leftrightarrow 3)]. \end{aligned} \quad (26)$$

Similarly, for  $\psi \rightarrow \rho_5(5^-) \pi \rightarrow \pi^+ \pi^- \pi^0$ , the amplitude should be

$$\begin{aligned} U_{\rho_5}^\mu &= (\mathbf{I}_1 \times \mathbf{I}_2) \cdot \mathbf{I}_3 \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma \tilde{T}_{(\rho_5 3)}^{(5)\nu\alpha\beta\gamma\delta} \tilde{t}_{(12)\alpha\beta\gamma\delta}^{(5)\lambda} f_{(12)}^{(\rho_5)} + (1 \leftrightarrow 3) + (2 \leftrightarrow 3) \\ &= -i \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma [\tilde{T}_{(\rho_5 3)}^{(5)\nu\alpha\beta\gamma\delta} \tilde{t}_{(12)\alpha\beta\gamma\delta}^{(5)\lambda} f_{(12)}^{(\rho_5)} - (1 \leftrightarrow 3) - (2 \leftrightarrow 3)]. \end{aligned} \quad (27)$$

If one considers a small isospin symmetry breaking effect, a free parameter can be multiplied into the term corresponding to the  $\rho^0$  intermediate state.

## 2.2 $\psi \rightarrow K^+ K^- \pi^0$

This channel is similar to  $\pi^+ \pi^- \pi^0$ . However, we now need to consider resonances for both  $K\pi$  and  $K^+ K^-$  subsystems. Numbering  $K^+$ ,  $K^-$ ,  $\pi^0$  as particle 1, 2, 3, the possible partial wave amplitudes are the following:

$$U_{\rho'}^\mu = \epsilon_{\mu\nu\lambda\sigma} p_1^\nu p_2^\lambda p_3^\sigma B_1(Q_{\psi\rho'3}) f_{(12)}^{(\rho')} B_1(Q_{\rho'12}), \quad (28)$$

$$U_{\rho_3}^\mu = \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma \tilde{T}_{(\rho_3 3)}^{(3)\nu\alpha\beta} \tilde{t}_{(12)\alpha\beta}^{(3)\lambda} \cdot f_{(12)}^{(\rho_3)}, \quad (29)$$

$$U_{\rho_5}^\mu = \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma \tilde{T}_{(\rho_5 3)}^{(5)\nu\alpha\beta\gamma\delta} \tilde{t}_{(12)\alpha\beta\gamma\delta}^{(5)\lambda} f_{(12)}^{(\rho_5)}, \quad (30)$$

$$U_{K^*}^\mu = \epsilon_{\mu\nu\lambda\sigma} p_1^\nu p_2^\lambda p_3^\sigma [B_1(Q_{\psi K^*2}) f_{(13)}^{(K^*)} B_1(Q_{K^*13}) + B_1(Q_{\psi K^*1}) f_{(23)}^{(K^*)} B_1(Q_{K^*23})], \quad (31)$$

$$U_{K_3^*}^\mu = \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma [\tilde{T}_{(K_3^* 2)}^{(3)\nu\alpha\beta} \tilde{t}_{(13)\alpha\beta}^{(3)\lambda} \cdot f_{(13)}^{(K_3^*)} - (1 \leftrightarrow 2)], \quad (32)$$

$$U_{K_5^*}^\mu = \epsilon_{\mu\nu\lambda\sigma} p_\psi^\sigma [\tilde{T}_{(K_5^* 2)}^{(5)\nu\alpha\beta\gamma\delta} \tilde{t}_{(13)\alpha\beta\gamma\delta}^{(5)\lambda} \cdot f_{(13)}^{(K_5^*)} - (1 \leftrightarrow 2)]. \quad (33)$$

## 2.3 $\psi \rightarrow \phi \pi^+ \pi^- \rightarrow K^+ K^- \pi^+ \pi^-$

For this channel,  $\phi$  is reconstructed from two kaons; most possible intermediate states are  $\phi$  plus an isospin zero resonance,  $f_0$  or  $f_2$ , which decays into two pions. The  $f_4$  is unlikely to be produced, because  $\psi$  mass is not far from  $\phi f_4$  threshold and the decay requires  $L = 2$  between  $\phi$  and  $f_4$ , hence a strong centrifugal barrier. For  $\psi \rightarrow \phi f_J$  in an orbital angular momentum  $L$  state, the conservation of total angular momentum requires

$$\mathbf{S}_\psi = \mathbf{S} + \mathbf{L} \quad (34)$$

where

$$\mathbf{S} = \mathbf{S}_\phi + \mathbf{J}. \quad (35)$$

In the following, we use notation  $\langle \phi f_J | LS \rangle$  to denote the corresponding partial wave amplitude  $U_i^\mu$ . We number the  $K^+$ ,  $K^-$ ,  $\pi^+$ ,  $\pi^-$  as particle 1, 2, 3, 4, respectively. Then we have two independent partial wave amplitudes for each  $f_0$  production. In the general formalism, they may be written:

$$\langle \phi f_0 | 01 \rangle = \tilde{t}_{(12)}^{(1)\mu} f_{(12)}^{(\phi)} f_{(34)}^{(f_0)}, \quad (36)$$

$$\langle \phi f_0 | 21 \rangle = \tilde{T}_{(\phi f_0)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(34)}^{(f_0)}. \quad (37)$$

For the very narrow  $\phi$  resonance, the  $\ell = 1$  centrifugal barrier factor for  $\phi$  decay has negligible effect on the  $\phi$  line-shape and can be dropped. The expression for  $t_{(12)}^{(1)\mu}$  simplifies to

$$t_{(12)}^{(1)\mu} = \tilde{r}^\mu = q_{(12)}^\mu.$$

In the last step, we use the fact that  $K^+$  and  $K^-$  have equal masses. Then Eqs. (36) and (37) become:

$$\langle \phi f_0 | 01 \rangle = q_{(12)}^\mu f_{(12)}^{(\phi)} f_{(34)}^{(f_0)}, \quad (38)$$

$$\langle \phi f_0 | 21 \rangle = \tau^{\mu\nu} q_{(12)\nu} B_2(Q_{\psi\phi f_0}) f_{(12)}^{(\phi)} f_{(34)}^{(f_0)}, \quad (39)$$

where  $\tau^{\mu\nu}$  is the  $L = 2$  operator

$$\tau^{\mu\nu} = q_{(12)}^\mu q_{(12)}^\nu - \frac{1}{3}(q_{(12)} \cdot q_{(12)})g^{\mu\nu}. \quad (40)$$

For each  $f_2$  production, there are five independent partial waves, which we retain in their general form:

$$\langle \phi f_2 | 01 \rangle = \tilde{t}_{(34)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(34)}^{(f_2)}, \quad (41)$$

$$\langle \phi f_2 | 21 \rangle = \tilde{T}_{(\phi f_2)}^{(2)\mu\alpha} \tilde{t}_{(34)\alpha\nu}^{(2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(34)}^{(f_2)}, \quad (42)$$

$$\langle \phi f_2 | 22 \rangle = \epsilon^{\mu\alpha\beta\gamma} p_{\psi\alpha} \tilde{T}_{(\phi f_2)\beta}^{(2)\delta} [\epsilon_{\gamma\lambda\sigma\nu} \tilde{t}_{(34)\delta}^{(2)\lambda} + \epsilon_{\delta\lambda\sigma\nu} \tilde{t}_{(34)\gamma}^{(2)\lambda}] p_{\psi}^\sigma \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(34)}^{(f_2)}, \quad (43)$$

$$\langle \phi f_2 | 23 \rangle = P^{(3)\mu\alpha\beta\gamma\delta\nu} (p_\psi) \tilde{T}_{(\phi f_2)\alpha\beta}^{(2)} \tilde{t}_{(34)\gamma\delta}^{(2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(34)}^{(f_2)}, \quad (44)$$

$$\langle \phi f_2 | 43 \rangle = \tilde{T}_{(\phi f_2)}^{(4)\mu\lambda\sigma} \tilde{t}_{(12)\nu}^{(1)} \tilde{t}_{(34)\lambda\sigma}^{(2)} f_{(12)}^{(\phi)} f_{(34)}^{(f_2)}. \quad (45)$$

There is no established resonance decaying into  $\phi\pi$ . However, there are speculations about  $(s\bar{s}q\bar{q})$  four-quark states which could decay to  $\phi\pi$ . So here we also give some partial wave amplitudes for  $\psi \rightarrow X\pi$  with the intermediate resonance  $X$  further decaying to  $\phi\pi$ . For  $X$  being a  $\rho'(1^{--})$  state, there is only one independent amplitude since both  $\psi \rightarrow \rho'\pi$  and  $\rho' \rightarrow \phi\pi$  are limited to a  $P$  wave.

$$U_{\rho'}^\mu = \epsilon_{\alpha\beta\gamma}^\mu p_{\psi}^\alpha [\tilde{T}_{(\rho'3)}^{(1)\beta} \epsilon^{\gamma\delta\sigma\lambda} p_{\psi\delta} \tilde{t}_{(\phi 4)\sigma}^{(1)} \tilde{t}_{(12)\lambda}^{(1)} f_{(12)}^{(\phi)} f_{(\phi 4)}^{(\rho')} + \tilde{T}_{(\rho'4)}^{(1)\beta} \epsilon^{\gamma\delta\sigma\lambda} p_{\psi\delta} \tilde{t}_{(\phi 3)\sigma}^{(1)} \tilde{t}_{(12)\lambda}^{(1)} f_{(12)}^{(\phi)} f_{(\phi 3)}^{(\rho')}] \quad (46)$$

For  $X$  being a  $b_1(1^{+-})$  state, there are four independent amplitudes since both  $\psi \rightarrow b_1\pi$  and  $b_1 \rightarrow \phi\pi$  can have both S and D waves.

$$U_{b_1 SS}^\mu = \tilde{g}_{(123)}^{\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(123)}^{(b_1)} + \tilde{g}_{(124)}^{\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(124)}^{(b_1)}, \quad (47)$$

$$U_{b_1 SD}^\mu = \tilde{t}_{(\phi 3)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(123)}^{(b_1)} + \tilde{t}_{(\phi 4)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(124)}^{(b_1)}, \quad (48)$$

$$U_{b_1 DS}^\mu = \tilde{T}_{(b_1 4)}^{(2)\mu\lambda} \tilde{g}_{(123)\lambda\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(123)}^{(b_1)} + \tilde{T}_{(b_1 3)}^{(2)\mu\lambda} \tilde{g}_{(124)\lambda\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(124)}^{(b_1)}, \quad (49)$$

$$U_{b_1 DD}^\mu = \tilde{T}_{(b_1 4)}^{(2)\mu\lambda} \tilde{t}_{(\phi 3)\lambda\nu}^{(2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(123)}^{(b_1)} + \tilde{T}_{(b_1 3)}^{(2)\mu\lambda} \tilde{t}_{(\phi 4)\lambda\nu}^{(2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(124)}^{(b_1)}. \quad (50)$$

## 2.4 $\psi \rightarrow \omega K^+ K^- \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-$

The formulae for this channel are quite similar to those in the previous subsection for the  $\psi \rightarrow \phi \pi^+ \pi^-$ . If we number the  $\pi^0, \pi^+, \pi^-, K^+, K^-$  as 0, 1, 2, 3, 4, then we can get corresponding partial wave amplitudes by simply replacing  $\tilde{t}_{(12)}^{(1)\mu}$  in equations of the previous subsection by  $\omega^\mu$  defined as

$$\omega^\mu = \epsilon_{\nu\lambda\sigma}^\mu p_1^\nu p_2^\lambda p_0^\sigma \left[ B_1(Q_{\omega\rho 0}) f_{(12)}^{(\rho)} B_1(Q_{\rho 12}) + B_1(Q_{\omega\rho 2}) f_{(10)}^{(\rho)} B_1(Q_{\rho 10}) + B_1(Q_{\omega\rho 1}) f_{(20)}^{(\rho)} B_1(Q_{\rho 20}) \right] \quad (51)$$

and replacing  $f_{(12)}^{(\phi)}$  by  $f_{(012)}^{(\omega)}$ .

## 2.5 $\psi \rightarrow K^+ \pi^- K^- \pi^+$

We label  $K^+, \pi^-, K^-, \pi^+$  as 1,2,3,4. For  $\rho a_0$  and  $\rho a_2$  intermediate states, the formulae are the same as for  $\phi f_0$  and  $\phi f_2$  intermediate states with a trivial exchange between pions and kaons. For  $KKK^* \rightarrow (KK^*\pi \text{ or } K\rho K)$ , or  $\pi\rho' \rightarrow K^*K$  intermediate states, the formulae are the same as for the  $\pi\rho' \rightarrow \pi\phi\pi$  intermediate state with proper recombination of particles. For  $KK_1^* \rightarrow KK^*\pi \text{ or } K\rho K$  intermediate states, the formulae are the same as for the  $\pi b_1 \rightarrow \pi\phi\pi$  intermediate state with proper recombination of particles. So all the formulae given in the subsection on  $\phi\pi^+\pi^-$  may be applied here. In addition, there are many more possible intermediate states. We list additional formulae for some obvious large intermediate states. Note for a resonance with the negative C-parity, it decays to  $K_{j_1}^* \bar{K}_{j_2}^*$  with a relative minus sign to its charge conjugate state  $\bar{K}_{j_1}^* K_{j_2}^*$ .

$$\langle K^* K_0^* | 01 \rangle = \tilde{t}_{(12)}^{(1)\mu} f_{(12)}^{(K^*)} f_{(34)}^{(\bar{K}_0^*)} - \tilde{t}_{(34)}^{(1)\mu} f_{(34)}^{(\bar{K}^*)} f_{(12)}^{(K_0^*)}, \quad (52)$$

$$\langle K^* K_0^* | 21 \rangle = \tilde{T}_{((12)(34))}^{(2)\mu\nu} [\tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(K^*)} f_{(34)}^{(K_0^*)} - \tilde{t}_{(34)\nu}^{(1)} f_{(34)}^{(K^*)} f_{(12)}^{(K_0^*)}], \quad (53)$$

$$\langle K^* K_2^* | 01 \rangle = \tilde{t}_{(34)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(K^*)} f_{(34)}^{(K_2^*)} - \tilde{t}_{(12)}^{(2)\mu\nu} \tilde{t}_{(34)\nu}^{(1)} f_{(34)}^{(K^*)} f_{(12)}^{(K_2^*)}, \quad (54)$$

$$\langle K^* K_2^* | 21 \rangle = \tilde{T}_{((12)(34))}^{(2)\mu\alpha} [\tilde{t}_{(34)\alpha\nu}^{(2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(K^*)} f_{(34)}^{(K_2^*)} - \tilde{t}_{(12)\alpha\nu}^{(2)} \tilde{t}_{(34)\nu}^{(1)} f_{(34)}^{(K^*)} f_{(12)}^{(K_2^*)}], \quad (55)$$

$$\begin{aligned} \langle K^* K_2^* | 22 \rangle &= \epsilon^{\mu\alpha\beta\gamma} p_{\psi\alpha} \tilde{T}_{((12)(34))\beta}^{(2)\delta} \{ [\epsilon_{\gamma\lambda\sigma\nu} \tilde{t}_{(34)\delta}^{(2)\lambda} + \epsilon_{\delta\lambda\sigma\nu} \tilde{t}_{(34)\gamma}^{(2)\lambda}] p_\psi^\sigma \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(K^*)} f_{(34)}^{(K_2^*)} \\ &\quad - [\epsilon_{\gamma\lambda\sigma\nu} \tilde{t}_{(12)\delta}^{(2)\lambda} + \epsilon_{\delta\lambda\sigma\nu} \tilde{t}_{(12)\gamma}^{(2)\lambda}] p_\psi^\sigma \tilde{t}_{(34)\nu}^{(1)} f_{(34)}^{(K^*)} f_{(12)}^{(K_2^*)} \}, \end{aligned} \quad (56)$$

$$\langle K^* K_2^* | 23 \rangle = P^{(3)\mu\alpha\beta\gamma\delta\nu} (p_\psi) \tilde{T}_{((12)(34))\alpha\beta}^{(2)} [\tilde{t}_{(34)\gamma\delta}^{(2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(K^*)} f_{(34)}^{(K_2^*)} - \tilde{t}_{(12)\gamma\delta}^{(2)} \tilde{t}_{(34)\nu}^{(1)} f_{(34)}^{(K^*)} f_{(12)}^{(K_2^*)}], \quad (57)$$

$$\langle K^* K_2^* | 43 \rangle = \tilde{T}_{((12)(34))}^{(4)\mu\nu\lambda\sigma} [\tilde{t}_{(12)\nu}^{(1)} \tilde{t}_{(34)\lambda\sigma}^{(2)} f_{(12)}^{(K^*)} f_{(34)}^{(K_2^*)} - \tilde{t}_{(34)\nu}^{(1)} \tilde{t}_{(12)\lambda\sigma}^{(2)} f_{(34)}^{(K^*)} f_{(12)}^{(K_2^*)}], \quad (58)$$

$$\langle K_0^* K_0^* | 10 \rangle = \tilde{T}_{((12)(34))}^{(1)\mu} [f_{(12)}^{K_0^*} f_{(34)}^{K_0^*} + f_{(34)}^{K_0^*} f_{(12)}^{K_0^*}], \quad (59)$$

$$\langle K_0^* K_2^* | 12 \rangle = \tilde{T}_{((12)(34))\nu}^{(1)} [\tilde{t}_{(34)}^{(2)\mu\nu} f_{(12)}^{(K_0^*)} f_{(34)}^{(K_2^*)} + \tilde{t}_{(12)}^{(2)\mu\nu} f_{(34)}^{(K_0^*)} f_{(12)}^{(K_2^*)}], \quad (60)$$



$$\langle K_0^* K_2^* | 32 \rangle = \tilde{T}_{((12)(34))}^{(3)\mu\nu\lambda} [\tilde{t}_{(34)\nu\lambda}^{(2)} f_{(12)}^{(K_0^*)} f_{(34)}^{(K_2^*)} + \tilde{t}_{(12)\nu\lambda}^{(2)} f_{(34)}^{(K_0^*)} f_{(12)}^{(K_2^*)}], \quad (61)$$

$$\langle K^* K'^* | 10 \rangle = \tilde{T}_{((12)(34))}^{(1)\mu} \tilde{t}_{(12)}^{(1)\alpha} \tilde{t}_{(34)\alpha}^{(1)} [f_{(12)}^{K^*} f_{(34)}^{K'^*} + f_{(34)}^{K^*} f_{(12)}^{K'^*}], \quad (62)$$

$$\langle K^* K'^* | 11 \rangle = \epsilon^{\mu\alpha\beta\gamma} p_{\psi\alpha} \epsilon_{\beta\nu\lambda\sigma} p_{\psi\nu} \tilde{t}_{(12)}^{(1)\lambda} \tilde{t}_{(34)}^{(1)\sigma} \tilde{T}_{((12)(34))\gamma}^{(1)} [f_{(12)}^{K^*} f_{(34)}^{K'^*} - f_{(34)}^{K^*} f_{(12)}^{K'^*}], \quad (63)$$

$$\langle K^* K'^* | 12 \rangle = P^{(2)\mu\nu\alpha\beta} (p_{\psi}) \tilde{t}_{(12)\alpha}^{(1)} \tilde{t}_{(34)\beta}^{(1)} \tilde{T}_{((12)(34))\nu}^{(1)} [f_{(12)}^{K^*} f_{(34)}^{K'^*} + f_{(34)}^{K^*} f_{(12)}^{K'^*}], \quad (64)$$

$$\langle K^* K'^* | 32 \rangle = \tilde{T}_{((12)(34))}^{(3)\mu\nu\lambda} \tilde{t}_{(12)\nu}^{(1)} \tilde{t}_{(34)\lambda}^{(1)} [f_{(12)}^{K^*} f_{(34)}^{K'^*} + f_{(34)}^{K^*} f_{(12)}^{K'^*}]. \quad (65)$$

Smaller contribution from  $KK_2^*$  with  $K_2^* \rightarrow K^*\pi$  or  $K\rho$  may needed. The corresponding formulae are

$$\begin{aligned} \langle KK_2^*(K^*\pi) | 22 \rangle &= \epsilon^{\mu\nu\lambda\sigma} p_{\psi\sigma} [\tilde{T}_{(K_2^*3)}^{(2)\delta} P_{\lambda\delta\alpha\beta}^{(2)}(p_{(124)}) \epsilon^{\alpha\gamma\eta\omega} p_{(124)\gamma} \tilde{t}_{(K^*4)\eta}^{(2)\beta} \tilde{t}_{(12)\omega}^{(1)} f_{(124)}^{(K_2^*)} f_{(12)}^{(K^*)} \\ &\quad - \tilde{T}_{(K_2^*1)\nu}^{(2)\delta} P_{\lambda\delta\alpha\beta}^{(2)}(p_{(234)}) \epsilon^{\alpha\gamma\eta\omega} p_{(234)\gamma} \tilde{t}_{(K^*2)\eta}^{(2)\beta} \tilde{t}_{(34)\omega}^{(1)}] f_{(234)}^{(K_2^*)} f_{(34)}^{(K^*)}, \quad (66) \end{aligned}$$

$$\begin{aligned} \langle KK_{2^+}^*(K\rho) | 22 \rangle &= \epsilon^{\mu\nu\lambda\sigma} p_{\psi\sigma} [\tilde{T}_{(K_2^*3)}^{(2)\delta} P_{\lambda\delta\alpha\beta}^{(2)}(p_{(124)}) \epsilon^{\alpha\gamma\eta\omega} p_{(124)\gamma} \tilde{t}_{(\rho1)\eta}^{(2)\beta} \tilde{t}_{(24)\omega}^{(1)} f_{(124)}^{(K_2^*)} f_{(24)}^{(\rho)} \\ &\quad - \tilde{T}_{(K_2^*1)\nu}^{(2)\delta} P_{\lambda\delta\alpha\beta}^{(2)}(p_{(234)}) \epsilon^{\alpha\gamma\eta\omega} p_{(234)\gamma} \tilde{t}_{(\rho3)\eta}^{(2)\beta} \tilde{t}_{(24)\omega}^{(1)}] f_{(234)}^{(K_2^*)} f_{(24)}^{(\rho)}. \quad (67) \end{aligned}$$

Some other intermediate states may also need to be considered. The formulae for  $\psi \rightarrow KK_{2^-}^*$  with  $K_{2^-}^* \rightarrow K_2^*\pi$  or  $K^*\pi$  or  $\rho K$  are

$$\begin{aligned} \langle KK_{2^-}^*(K_2^*\pi) | 12 \rangle &= P^{(2)\mu\nu\lambda\sigma} (p_{(124)}) \tilde{T}_{(K_2^*3)\nu}^{(1)} P_{\lambda\sigma\alpha\beta}^{(2)}(p_{(12)}) \tilde{t}_{(12)}^{(2)\alpha\beta} f_{(124)}^{(K_2^*-)} f_{(12)}^{(K_2^*)} \\ &\quad - P^{(2)\mu\nu\lambda\sigma} (p_{(234)}) \tilde{T}_{(K_2^*1)\nu}^{(1)} P_{\lambda\sigma\alpha\beta}^{(2)}(p_{(34)}) \tilde{t}_{(34)}^{(2)\alpha\beta} f_{(234)}^{(K_2^*-)} f_{(34)}^{(K_2^*)}, \quad (68) \end{aligned}$$

$$\begin{aligned} \langle KK_{2^-}^*(K_2^*\pi) | 32 \rangle &= \tilde{T}_{(K_2^*3)}^{(3)\mu\nu\gamma} P_{\nu\gamma\lambda\sigma}^{(2)}(p_{(124)}) P^{(2)\lambda\sigma\alpha\beta} (p_{(12)}) \tilde{t}_{(12)\alpha\beta}^{(2)} f_{(124)}^{(K_2^*-)} f_{(12)}^{(K_2^*)} \\ &\quad - \tilde{T}_{(K_2^*1)}^{(3)\mu\nu\gamma} P_{\nu\gamma\lambda\sigma}^{(2)}(p_{(234)}) P^{(2)\lambda\sigma\alpha\beta} (p_{(34)}) \tilde{t}_{(34)\alpha\beta}^{(2)} f_{(234)}^{(K_2^*-)} f_{(34)}^{(K_2^*)}, \quad (69) \end{aligned}$$

$$\begin{aligned} \langle KK_{2^-}^*(K^*\pi) | 12 \rangle &= P^{(2)\mu\nu\lambda\sigma} (p_{(124)}) \tilde{T}_{(K_2^*3)\nu}^{(1)} P_{\lambda\sigma\alpha\beta}^{(2)}(p_{(12)}) \tilde{t}_{(K^*4)\alpha}^{(1)} \tilde{t}_{(12)\beta}^{(1)} f_{(124)}^{(K_2^*-)} f_{(12)}^{(K^*)} \\ &\quad - P^{(2)\mu\nu\lambda\sigma} (p_{(234)}) \tilde{T}_{(K_2^*1)\nu}^{(1)} P_{\lambda\sigma\alpha\beta}^{(2)}(p_{(34)}) \tilde{t}_{(K^*2)\alpha}^{(1)} \tilde{t}_{(34)\beta}^{(1)} f_{(234)}^{(K_2^*-)} f_{(34)}^{(K^*)}, \quad (70) \end{aligned}$$

$$\begin{aligned} \langle KK_{2^-}^*(K^*\pi) | 32 \rangle &= \tilde{T}_{(K_2^*3)}^{(3)\mu\nu\gamma} P_{\nu\gamma\lambda\sigma}^{(2)}(p_{(124)}) P^{(2)\lambda\sigma\alpha\beta} (p_{(12)}) \tilde{t}_{(K^*4)\alpha}^{(1)} \tilde{t}_{(12)\beta}^{(1)} f_{(124)}^{(K_2^*-)} f_{(12)}^{(K^*)} \\ &\quad - \tilde{T}_{(K_2^*1)}^{(3)\mu\nu\gamma} P_{\nu\gamma\lambda\sigma}^{(2)}(p_{(234)}) P^{(2)\lambda\sigma\alpha\beta} (p_{(34)}) \tilde{t}_{(K^*2)\alpha}^{(1)} \tilde{t}_{(34)\beta}^{(1)} f_{(234)}^{(K_2^*-)} f_{(34)}^{(K^*)} \quad (71) \end{aligned}$$

$$\begin{aligned} \langle KK_{2^-}^*(\rho K) | 12 \rangle &= P^{(2)\mu\nu\lambda\sigma} (p_{(124)}) \tilde{T}_{(K_2^*3)\nu}^{(1)} P_{\lambda\sigma\alpha\beta}^{(2)}(p_{(12)}) \tilde{t}_{(\rho1)\alpha}^{(1)} \tilde{t}_{(24)\beta}^{(1)} f_{(124)}^{(K_2^*-)} f_{(24)}^{(\rho)} \\ &\quad - P^{(2)\mu\nu\lambda\sigma} (p_{(234)}) \tilde{T}_{(K_2^*1)\nu}^{(1)} P_{\lambda\sigma\alpha\beta}^{(2)}(p_{(34)}) \tilde{t}_{(\rho3)\alpha}^{(1)} \tilde{t}_{(24)\beta}^{(1)} f_{(234)}^{(K_2^*-)} f_{(24)}^{(\rho)}, \quad (72) \end{aligned}$$

$$\langle KK_{2^-}^*(\rho K) | 32 \rangle = \tilde{T}_{(K_2^*3)}^{(3)\mu\nu\gamma} P_{\nu\gamma\lambda\sigma}^{(2)}(p_{(124)}) P^{(2)\lambda\sigma\alpha\beta} (p_{(12)}) \tilde{t}_{(\rho1)\alpha}^{(1)} \tilde{t}_{(24)\beta}^{(1)} f_{(124)}^{(K_2^*-)} f_{(24)}^{(\rho)}$$

$$-\tilde{T}_{(\bar{K}_2^*-1)}^{(3)\mu\nu\gamma} P_{\nu\gamma\lambda\sigma}^{(2)}(p_{(234)}) P^{(2)\lambda\sigma\alpha\beta}(p_{(34)}) \tilde{t}_{(\rho 3)\alpha}^{(1)} \tilde{t}_{(24)\beta}^{(1)} f_{(234)}^{(K_2^*)} f_{(24)}^{(\rho)}. \quad (73)$$

The formulae for  $\psi \rightarrow KK'$  with  $K'$  the excited  $0^-$  state and  $K' \rightarrow K^*\pi$  or  $\rho K$  or  $K_0^*\pi$  are

$$\langle KK'(K^*\pi)|10\rangle = \tilde{T}_{(K'3)}^{(1)\mu} \tilde{t}_{K^*4}^{(1)\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(124)}^{(K')} f_{(12)}^{(K^*)} - \tilde{T}_{(\bar{K}'1)}^{(1)\mu} \tilde{t}_{\bar{K}^*2}^{(1)\nu} \tilde{t}_{(34)\nu}^{(1)} f_{(234)}^{(K')} f_{(34)}^{(K^*)}, \quad (74)$$

$$\langle KK'(\rho K)|10\rangle = \tilde{T}_{(K'3)}^{(1)\mu} \tilde{t}_{\rho 1}^{(1)\nu} \tilde{t}_{(24)\nu}^{(1)} f_{(124)}^{(K')} f_{(24)}^{(\rho)} - \tilde{T}_{(\bar{K}'1)}^{(1)\mu} \tilde{t}_{\rho 3}^{(1)\nu} \tilde{t}_{(24)\nu}^{(1)} f_{(234)}^{(K')} f_{(24)}^{(\rho)}, \quad (75)$$

$$\langle KK'(K_0^*\pi)|10\rangle = \tilde{T}_{(K'3)}^{(1)\mu} f_{(124)}^{(K')} f_{(12)}^{(K_0^*)} - \tilde{T}_{(\bar{K}'1)}^{(1)\mu} f_{(234)}^{(K')} f_{(34)}^{(K_0^*)}. \quad (76)$$

## 2.6 $\psi \rightarrow \phi\pi^+\pi^-\pi^+\pi^- \rightarrow K^+K^-\pi^+\pi^-\pi^+\pi^-$

As for the  $\psi \rightarrow \phi\pi^+\pi^-$  channel, the dominant intermediate states are also  $\phi f_0$  and  $\phi f_2$ . The  $f_0$  resonances decay to  $\pi^+\pi^-\pi^+\pi^-$  usually through  $\sigma\sigma$  and  $\rho\rho$ ; and  $f_2$  resonances decay to  $\pi^+\pi^-\pi^+\pi^-$  usually through  $\sigma\sigma$ ,  $\rho\rho$  and  $f_2(1270)\sigma$ . We assume a similar notation to the  $\psi \rightarrow \phi\pi^+\pi^-$  case and number the additional  $\pi^+\pi^-$  as particle 5, 6. Then the corresponding partial wave amplitudes involving  $f_J \rightarrow \sigma\sigma$  are:

$$\langle \phi f_0|01\rangle_{(\sigma\sigma)} = \tilde{t}_{(12)}^{(1)\mu} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_0)} [f_{(34)}^{(\sigma)} f_{(56)}^{(\sigma)} + f_{(36)}^{(\sigma)} f_{(45)}^{(\sigma)}], \quad (77)$$

$$\langle \phi f_0|21\rangle_{(\sigma\sigma)} = \tilde{T}_{(\phi f_0)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_0)} [f_{(34)}^{(\sigma)} f_{(56)}^{(\sigma)} + f_{(36)}^{(\sigma)} f_{(45)}^{(\sigma)}], \quad (78)$$

$$\langle \phi f_2|01\rangle_{(\sigma\sigma)} = T_{(\sigma\sigma)}^{(f_2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_2)}, \quad (79)$$

$$\langle \phi f_2|21\rangle_{(\sigma\sigma)} = \tilde{T}_{(\phi f_2)}^{(2)\mu\alpha} \tilde{T}_{(\sigma\sigma)\alpha\nu}^{(f_2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_2)}, \quad (80)$$

$$\langle \phi f_2|22\rangle_{(\sigma\sigma)} = \epsilon^{\mu\alpha\beta\gamma} p_{\psi\alpha} \tilde{T}_{(\phi f_2)\beta}^{(2)\delta} [\epsilon_{\gamma\lambda\sigma\nu} \tilde{T}_{(\sigma\sigma)\delta}^{(f_2)\lambda} + \epsilon_{\delta\lambda\sigma\nu} \tilde{T}_{(\sigma\sigma)\gamma}^{(f_2)\lambda}] p_{\psi}^{\sigma} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_2)}, \quad (81)$$

$$\langle \phi f_2|23\rangle_{(\sigma\sigma)} = P^{(3)\mu\alpha\beta\gamma\delta\nu}(p_{\psi}) \tilde{T}_{(\phi f_2)\alpha\beta}^{(2)} \tilde{T}_{(\sigma\sigma)\gamma\delta}^{(f_2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_2)}, \quad (82)$$

$$\langle \phi f_2|43\rangle_{(\sigma\sigma)} = \tilde{T}_{(\phi f_2)}^{(4)\mu\nu\lambda\sigma} \tilde{t}_{(12)\nu}^{(1)} T_{(\sigma\sigma)\lambda\sigma}^{(f_2)} f_{(12)}^{(\phi)} f_{(\sigma\sigma)}^{(f_2)} \quad (83)$$

with

$$T_{(\sigma\sigma)}^{(f_2)\mu\nu} = \tilde{t}_{(\sigma_{34}\sigma_{56})}^{(2)\mu\nu} f_{(34)}^{(\sigma)} f_{(56)}^{(\sigma)} + \tilde{t}_{(\sigma_{36}\sigma_{45})}^{(2)\mu\nu} f_{(36)}^{(\sigma)} f_{(45)}^{(\sigma)}. \quad (84)$$

For  $f_J \rightarrow \rho\rho$ , if we limit  $\rho\rho$  to a relative  $l=0$  state, then the corresponding partial wave amplitudes are:

$$\langle \phi f_0|01\rangle_{(\rho\rho)} = \tilde{t}_{(12)}^{(1)\mu} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_0)} [f_{(34)}^{(\rho)} f_{(56)}^{(\rho)} \tilde{t}_{(34)}^{(1)\alpha\beta} \tilde{t}_{(56)\alpha\beta}^{(1)} + f_{(36)}^{(\rho)} f_{(45)}^{(\rho)} \tilde{t}_{(36)}^{(1)\alpha\beta} \tilde{t}_{(45)\alpha\beta}^{(1)}], \quad (85)$$

$$\langle \phi f_0|21\rangle_{(\rho\rho)} = \tilde{T}_{(\phi f_0)}^{(2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_0)} [f_{(34)}^{(\rho)} f_{(56)}^{(\rho)} \tilde{t}_{(34)}^{(1)\alpha\beta} \tilde{t}_{(56)\alpha\beta}^{(1)} + f_{(36)}^{(\rho)} f_{(45)}^{(\rho)} \tilde{t}_{(36)}^{(1)\alpha\beta} \tilde{t}_{(45)\alpha\beta}^{(1)}], \quad (86)$$

$$\langle \phi f_2|01\rangle_{(\rho\rho)} = T_{(\rho\rho)}^{(f_2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_2)}, \quad (87)$$

$$\langle \phi f_2|21\rangle_{(\rho\rho)} = \tilde{T}_{(\phi f_2)}^{(2)\mu\alpha} \tilde{T}_{(\rho\rho)\alpha\nu}^{(f_2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_2)}, \quad (88)$$

$$\langle \phi f_2 | 22 \rangle_{(\rho\rho)} = \epsilon^{\mu\alpha\beta\gamma} p_{\psi\alpha} \tilde{T}_{(\phi f_2)\beta}^{(2)\delta} [\epsilon_{\gamma\lambda\sigma\nu} \tilde{T}_{(\rho\rho)\delta}^{(f_2)\lambda} + \epsilon_{\delta\lambda\sigma\nu} \tilde{T}_{(\rho\rho)\gamma}^{(f_2)\lambda}] p_{\psi}^{\sigma} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_2)}, \quad (89)$$

$$\langle \phi f_2 | 23 \rangle_{(\rho\rho)} = P^{(3)\mu\alpha\beta\gamma\delta\nu} (p_{\psi}) \tilde{T}_{(\phi f_2)\alpha\beta}^{(2)} \tilde{T}_{(\rho\rho)\gamma\delta}^{(f_2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_2)}, \quad (90)$$

$$\langle \phi f_2 | 43 \rangle_{(\rho\rho)} = \tilde{T}_{(\phi f_2)}^{(4)\mu\nu\lambda\sigma} \tilde{t}_{(12)\nu}^{(1)} T_{(\rho\rho)\lambda\sigma}^{(f_2)} f_{(12)}^{(\phi)} f_{(\rho\rho)}^{(f_2)} \quad (91)$$

where

$$T_{(\rho\rho)}^{(f_2)\mu\nu} = P^{(2)\mu\nu\alpha\beta} (p_{f_2}) \left[ \tilde{t}_{(34)\alpha}^{(1)} \tilde{t}_{(56)\beta}^{(1)} f_{(34)}^{(\rho)} f_{(56)}^{(\rho)} + \tilde{t}_{(36)\alpha}^{(1)} \tilde{t}_{(45)\beta}^{(1)} f_{(36)}^{(\rho)} f_{(45)}^{(\rho)} \right]. \quad (92)$$

For  $f_2 \rightarrow f_2(1270)\sigma$ , if we also limit  $f_2(1270)\sigma$  to the  $l = 0$  state, then we have their corresponding partial wave amplitudes:

$$\langle \phi f_2 | 01 \rangle_{(f_2\sigma)} = T_{(f_2\sigma)}^{(f_2)\mu\nu} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(f_2\sigma)}^{(f_2)}, \quad (93)$$

$$\langle \phi f_2 | 21 \rangle_{(f_2\sigma)} = \tilde{T}_{(\phi f_2)}^{(2)\mu\alpha} \tilde{T}_{(f_2\sigma)\alpha\nu}^{(f_2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(f_2\sigma)}^{(f_2)}, \quad (94)$$

$$\langle \phi f_2 | 22 \rangle_{(f_2\sigma)} = \epsilon^{\mu\alpha\beta\gamma} p_{\psi\alpha} \tilde{T}_{(\phi f_2)\beta}^{(2)\delta} [\epsilon_{\gamma\lambda\sigma\nu} \tilde{T}_{(f_2\sigma)\delta}^{(f_2)\lambda} + \epsilon_{\delta\lambda\sigma\nu} \tilde{T}_{(f_2\sigma)\gamma}^{(f_2)\lambda}] p_{\psi}^{\sigma} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(f_2\sigma)}^{(f_2)}, \quad (95)$$

$$\langle \phi f_2 | 23 \rangle_{(f_2\sigma)} = P^{(3)\mu\alpha\beta\gamma\delta\nu} (p_{\psi}) \tilde{T}_{(\phi f_2)\alpha\beta}^{(2)} \tilde{T}_{(f_2\sigma)\gamma\delta}^{(f_2)} \tilde{t}_{(12)\nu}^{(1)} f_{(12)}^{(\phi)} f_{(f_2\sigma)}^{(f_2)}, \quad (96)$$

$$\langle \phi f_2 | 43 \rangle_{(f_2\sigma)} = \tilde{T}_{(\phi f_2)}^{(4)\mu\nu\lambda\sigma} \tilde{t}_{(12)\nu}^{(1)} T_{(f_2\sigma)\lambda\sigma}^{(f_2)} f_{(12)}^{(\phi)} f_{(f_2\sigma)}^{(f_2)} \quad (97)$$

with

$$T_{(f_2\sigma)}^{(f_2)\mu\nu} = P^{(2)\mu\nu\alpha\beta} (p_{f_2}) \left[ \tilde{t}_{(34)\alpha\beta}^{(2)} f_{(34)}^{(f_2)} f_{(56)}^{(\sigma)} + \tilde{t}_{(56)\alpha\beta}^{(2)} f_{(56)}^{(f_2)} f_{(34)}^{(\sigma)} \right. \\ \left. + \tilde{t}_{(36)\alpha\beta}^{(2)} f_{(36)}^{(f_2)} f_{(45)}^{(\sigma)} + \tilde{t}_{(45)\alpha\beta}^{(2)} f_{(45)}^{(f_2)} f_{(36)}^{(\sigma)} \right]. \quad (98)$$

Unlike the  $\psi \rightarrow \phi\pi^+\pi^-$  channel, for  $\psi \rightarrow \phi\pi^+\pi^-\pi^+\pi^-$  it is possible to go through  $0^{--}$  resonances ( $\eta^*$ ) decaying to  $\pi^+\pi^-\pi^+\pi^-$  through  $\rho\rho$ . The corresponding partial wave is

$$U_{\eta^*}^{\mu} = \epsilon^{\mu\nu\alpha\beta} \tilde{t}_{(12)\nu}^{(1)} \tilde{T}_{(\phi\eta^*)\alpha}^{(1)} p_{\psi\beta} \epsilon_{\tau\sigma\gamma\eta} p_3^{\tau} p_4^{\sigma} p_5^{\gamma} p_6^{\eta} [f_{(34)}^{(\rho)} f_{(56)}^{(\rho)} B_1(Q_{\rho 34}) B_1(Q_{\rho 56}) B_1(Q_{\eta^* \rho 34 \rho 56}) \\ - f_{(36)}^{(\rho)} f_{(45)}^{(\rho)} B_1(Q_{\rho 36}) B_1(Q_{\rho 45}) B_1(Q_{\eta^* \rho 36 \rho 45})]. \quad (99)$$

Besides partial wave amplitudes given above, for  $\pi^+\pi^-\pi^+\pi^-$  final states, there are many other possible intermediate states, such as  $a_2\pi$ ,  $a_1\pi$ ,  $\pi(1300)\pi$  etc. Before performing partial wave analysis, one should check various invariant mass spectrum to see what resonances are present in the data and add the corresponding partial wave amplitudes.

### 3 Formalism for $\psi$ radiative decay to mesons

We denote the  $\psi$  polarization four-vector by  $\psi_{\mu}(m_1)$  and the polarization vector of the photon by  $e_{\nu}(m_2)$ . Then the general form for the decay amplitude is

$$A = \psi_{\mu}(m_1) e_{\nu}^*(m_2) A^{\mu\nu} = \psi_{\mu}(m_1) e_{\nu}^*(m_2) \sum_i \Lambda_i U_i^{\mu\nu}. \quad (100)$$

For the photon polarization four vector  $e_\nu$  with photon momentum  $q$ , there is the usual Lorentz orthogonality condition  $e_\nu q^\nu = 0$ . This is the same as for a massive vector meson. However, for the photon, there is an additional gauge invariance condition. Here we assume the Coulomb gauge in the  $\psi$  rest system, i.e.,  $e_\nu p_\psi^\nu = 0$ . Then we have [13]

$$\sum_m e_\mu^*(m) e_\nu(m) = -g_{\mu\nu} + \frac{q_\mu K_\nu + K_\mu q_\nu}{q \cdot K} - \frac{K \cdot K}{(q \cdot K)^2} q_\mu q_\nu \equiv -g_{\mu\nu}^{(\perp\perp)} \quad (101)$$

with  $K = p_\psi - q$  and  $e_\nu K^\nu = 0$ . The radiative decay cross section is:

$$\begin{aligned} \frac{d\sigma}{d\Phi_n} &= \frac{1}{2} \sum_{m_1=1}^2 \sum_{m_2=1}^2 \psi_\mu(m_1) e_\nu^*(m_2) A^{\mu\nu} \psi_{\mu'}^*(m_1) e_{\nu'}(m_2) A^{*\mu'\nu'} \\ &= -\frac{1}{2} \sum_{m_1=1}^2 \psi_\mu(m_1) \psi_{\mu'}^*(m_1) g_{\nu\nu'}^{(\perp\perp)} A^{\mu\nu} A^{*\mu'\nu'} \\ &= -\frac{1}{2} \sum_{\mu=1}^2 A_{\mu\nu} g_{\nu\nu'}^{(\perp\perp)} A^{*\mu\nu'} \\ &= -\frac{1}{2} \sum_{i,j} \Lambda_i \Lambda_j^* \sum_{\mu=1}^2 U_i^{\mu\nu} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'} \equiv \sum_{i,j} P_{ij} \cdot F_{ij} \end{aligned} \quad (102)$$

where

$$P_{ij} = P_{ji}^* = \Lambda_i \Lambda_j^*, \quad (103)$$

$$F_{ij} = F_{ji}^* = -\frac{1}{2} \sum_{\mu=1}^2 U_i^{\mu\nu} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'}. \quad (104)$$

Due to the special properties (massless and gauge invariance) of the photon, the number of independent partial wave amplitudes for a  $\psi$  radiative decay is smaller than for the corresponding decay to a massive vector meson. For example, for  $\psi \rightarrow \phi f_0$ , there are two independent partial wave amplitudes with orbital angular momentum  $L = 0$  and 2, respectively, which give different angular distributions; but for  $\psi \rightarrow \gamma f_0$ , with the gauge invariance condition, the two amplitudes will give the same angular distribution. So for the  $\psi$  radiative decay, the L-S scheme is not useful any more for choosing independent amplitudes. One may simply use momenta of the particles to construct covariant tensor amplitudes; it is sufficient to check the helicity amplitudes to make sure there is the right number of independent amplitudes. From the helicity formalism, it is easy to show that there is one independent amplitude for  $\psi$  radiative decay to a spin-0 meson, two independent amplitudes for  $\psi$  radiative decay to a spin-1 meson, and three independent amplitudes for  $\psi$  radiative decay to a meson with spin larger than 1.

### 3.1 $\psi$ radiative decay to two pseudoscalar mesons

We denote the two pseudoscalar mesons as  $\pi^+$  and  $\pi^-$ . For the decay vertex  $\psi \rightarrow \gamma f_J$ , there are two independent momenta which we choose to be  $p_\psi$  and the momentum of the photon  $q$ . We use these two momenta and spin wave functions of the three particles to construct the covariant tensor amplitudes.

For  $\psi \rightarrow \gamma f_0$ , the  $e_\mu$  can only contract with  $\psi^\mu$  since  $e_\mu p_\psi^\mu = e_\mu q^\mu = 0$ ; hence there is only one independent amplitude:

$$U_{\gamma f_0}^{\mu\nu} = g^{\mu\nu} f^{(f_0)}. \quad (105)$$

For  $\psi \rightarrow \gamma f_2$  or  $\psi \rightarrow \gamma f_4$ , the  $e_\mu$  may contract with  $\psi^\mu$  or with the spin wave function of  $f_J$ . Then  $\psi^\mu$  may contract with  $e_\mu$ , or  $q_\mu$ , or the spin wave function of  $f_J$ ; this gives three independent covariant tensor amplitudes for each  $f_J$ :

$$U_{(\gamma f_2)1}^{\mu\nu} = \tilde{t}^{(f_2)\mu\nu} f^{(f_2)}, \quad (106)$$

$$U_{(\gamma f_2)2}^{\mu\nu} = g^{\mu\nu} p_\psi^\alpha p_\psi^\beta \tilde{t}_{\alpha\beta}^{(f_2)} B_2(Q_{\Psi\gamma f_2}) f^{(f_2)}, \quad (107)$$

$$U_{(\gamma f_2)3}^{\mu\nu} = q^\mu \tilde{t}_\alpha^{(f_2)\nu} p_\psi^\alpha B_2(Q_{\psi\gamma f_2}) f^{(f_2)}, \quad (108)$$

$$U_{(\gamma f_4)1}^{\mu\nu} = \tilde{t}_{\alpha\beta}^{(f_4)\mu\nu} p_\psi^\alpha p_\psi^\beta B_2(Q_{\Psi\gamma f_4}) f^{(f_4)}, \quad (109)$$

$$U_{(\gamma f_4)2}^{\mu\nu} = g^{\mu\nu} \tilde{t}_{\alpha\beta\gamma\delta}^{(f_4)} p_\psi^\alpha p_\psi^\beta p_\psi^\gamma p_\psi^\delta B_4(Q_{\psi\gamma f_4}) f^{(f_4)}, \quad (110)$$

$$U_{(\gamma f_4)3}^{\mu\nu} = q^\mu \tilde{t}_{\alpha\beta\gamma}^{(f_4)\nu} p_\psi^\alpha p_\psi^\beta p_\psi^\gamma B_4(Q_{\Psi\gamma f_4}) f^{(f_4)} \quad (111)$$

where

$$\begin{aligned} \tilde{t}_{\mu_1 \dots \mu_J}^{(f_J)} &= \sum_m \phi_{\mu_1 \dots \mu_J}(p_{f_J}, m) \phi_{\mu'_1 \dots \mu'_J}(p_{f_J}, m) r_\pi^{\mu'_1} \dots r_\pi^{\mu'_J} B_J(Q_{f_J \pi \pi}) \\ &= P_{\mu_1 \dots \mu_J \mu'_1 \dots \mu'_J}(p_{f_J}) r_\pi^{\mu'_1} \dots r_\pi^{\mu'_J} B_J(Q_{f_J \pi \pi}) \end{aligned} \quad (112)$$

with  $J = 0, 2$ ; here  $r_\pi$  represents the relative momentum between two pseudoscalar mesons.

We use  $p_\psi$  instead of  $q$  to contract with  $\tilde{t}^{(f_J)}$  because  $q\tilde{t}^{(f_J)} = p_\psi\tilde{t}^{(f_J)}$  and  $p_\psi$  has only a time component in the  $\psi$  rest system. This makes the calculation simpler.

### 3.2 $\psi \rightarrow \gamma \eta \pi^+ \pi^-$

This is a three step process:  $\psi \rightarrow \gamma X$  with  $X \rightarrow yz$  and  $y \rightarrow \pi\pi$  or  $y \rightarrow \eta\pi$ . The amplitudes  $U_{\mu\nu}^i$  are listed using the notation:

$$\langle \gamma J^{PC} | (yz) i \rangle \quad (113)$$

where J,P,C are the intrinsic spin, parity and C-parity of the X particle, respectively. We denote  $\pi^+$ ,  $\pi^-$ ,  $\eta$  as 1,2,3, respectively. The possible  $J^{PC}$  for X are  $0^{-+}$ ,  $1^{++}$ ,  $1^{-+}$ ,  $2^{++}$ ,  $2^{-+}$ ,  $3^{++}$ ,  $3^{-+}$ , etc. For invariant mass below 2 GeV, we consider J up to 2. For  $\psi \rightarrow \gamma X$ , we choose two independent momenta  $p_\psi$  for  $\psi$  and  $q$  for the photon to be contracted with spin wave functions.

For the  $\psi \rightarrow \gamma 0^{-+}$  vertex, there is only one independent coupling,  $\epsilon_{\mu\nu\lambda\sigma} \psi^\mu e^\nu q^\lambda p_\psi^\sigma$ . With various possible  $yz$  states, we have  $U_{\mu\nu}^i$  for  $\psi \rightarrow \gamma 0^- \rightarrow \eta \pi^+ \pi^-$  as follows:

$$\langle \gamma 0^{-+} | (f_0 \eta) 1 \rangle = S_{\mu\nu} B_1(Q_{\psi\gamma X}) f_{(12)}^{(f_0)}, \quad (114)$$

$$\langle \gamma 0^{-+} | (a_0 \pi) 1 \rangle = S_{\mu\nu} B_1(Q_{\psi\gamma X}) (f_{(13)}^{(a_0)} + f_{(23)}^{(a_0)}), \quad (115)$$

$$\langle \gamma 0^{-+} | (f_2 \eta) 1 \rangle = S_{\mu\nu} B_1(Q_{\psi\gamma X}) f_{(12)}^{(f_2)} \tilde{t}_{(f_2\eta)\gamma\delta}^{(2)} \tilde{t}_{(12)}^{(2)\gamma\delta}, \quad (116)$$

$$\langle \gamma 0^{-+} | (a_2 \pi) 1 \rangle = S_{\mu\nu} B_1(Q_{\psi\gamma X}) \left\{ f_{(13)}^{(a_2)} \tilde{t}_{(a_2 2)\gamma\delta}^{(2)} \tilde{t}_{(13)}^{(2)\gamma\delta} + f_{(23)}^{(a_2)} \tilde{t}_{(a_2 1)\gamma\delta}^{(2)} \tilde{t}_{(23)}^{(2)\gamma\delta} \right\} \quad (117)$$

with  $S_{\mu\nu}$  defined as

$$S_{\mu\nu} = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha q^\beta. \quad (118)$$

For the  $\psi \rightarrow \gamma 1^{++}$  vertex, there are two independent couplings for each  $yz$ .

$$\langle \gamma 1^{++} | (f_0 \eta) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha \tilde{t}_{(\eta f_0)}^{(1)\beta} f_{(12)}^{(f_0)}, \quad (119)$$

$$\langle \gamma 1^{++} | (a_0 \pi) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha (\tilde{t}_{(a_0 1)}^{(1)\beta} f_{(23)}^{(a_0)} + \tilde{t}_{(a_0 2)}^{(1)\beta} f_{(13)}^{(a_0)}), \quad (120)$$

$$\langle \gamma 1^{++} | (f_0 \eta) 2 \rangle = q_\mu S_{\nu\beta} \tilde{t}_{(\eta f_0)}^{(1)\beta} B_2(Q_{\psi\gamma X}) f_{(12)}^{(f_0)}, \quad (121)$$

$$\langle \gamma 1^{++} | (a_0 \pi) 2 \rangle = q_\mu S_{\nu\beta} B_2(Q_{\psi\gamma X}) [t_{(a_0 1)}^{(1)\beta} f_{(23)}^{(a_0)} + t_{(a_0 2)}^{(1)\beta} f_{(13)}^{(a_0)}], \quad (122)$$

$$\langle \gamma 1^{++} | (f_2 \eta) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha \tilde{g}_X^{\beta\gamma} \tilde{t}_{(12)\gamma\delta}^{(2)} \tilde{t}_{(\eta f_2)}^{(1)\delta} f_{(12)}^{(f_2)}, \quad (123)$$

$$\langle \gamma 1^{++} | (a_2 \pi) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha \tilde{g}_X^{\beta\gamma} [\tilde{t}_{(13)\gamma\delta}^{(2)} \tilde{t}_{(a_2 2)}^{(1)\delta} f_{(13)}^{(a_2)} + \tilde{t}_{(23)\gamma\delta}^{(2)} \tilde{t}_{(a_2 1)}^{(1)\delta} f_{(23)}^{(a_2)}] \quad (124)$$

$$\langle \gamma 1^{++} | (f_2 \eta) 2 \rangle = q_\mu S_{\nu\beta} \tilde{g}_X^{\beta\beta'} \tilde{t}_{(12)\beta'\alpha'}^{(1)\alpha'} B_2(Q_{\psi\gamma X}) f_{(12)}^{(f_2)}, \quad (125)$$

$$\langle \gamma 1^{++} | (a_2 \pi) 2 \rangle = q_\mu S_{\nu\beta} \tilde{g}_X^{\beta\beta'} B_2(Q_{\psi\gamma X}) [\tilde{t}_{(13)\beta'\alpha'}^{(2)} \tilde{t}_{(a_2 2)}^{(1)\alpha'} f_{(13)}^{(a_2)} + \tilde{t}_{(23)\beta'\alpha'}^{(2)} \tilde{t}_{(a_2 1)}^{(1)\alpha'} f_{(23)}^{(a_2)}] \quad (126)$$

where  $\tilde{g}_X^{\alpha\beta} = g^{\alpha\beta} - \frac{p_X^\alpha p_X^\beta}{p_X^2}$ . For  $1^{++}$  decaying to  $f_2 \eta$  and  $a_2 \pi$ , the orbital angular momentum  $l$  could be 1 and 3; but we ignore the  $l = 3$  contribution because of the strong centrifugal barrier.

For  $\psi \rightarrow \gamma 1^{-+}$ , the exotic  $1^{-+}$  meson cannot decay into  $f_0 \eta$  and  $a_0 \pi$ . We have four  $U_{\mu\nu}^i$  amplitudes here:

$$\langle \gamma 1^{-+} | (f_2 \eta) 1 \rangle = g_{\mu\nu} S_{\gamma\delta} \tilde{t}_{(\eta f_2)}^{(2)\gamma\sigma} \tilde{t}_{(12)\sigma}^{(2)\delta} f_{(12)}^{(f_2)} B_1(Q_{\psi\gamma X}), \quad (127)$$

$$\langle \gamma 1^{-+} | (a_2 \pi) 1 \rangle = g_{\mu\nu} S_{\gamma\delta} [\tilde{t}_{(a_2 2)}^{(2)\gamma\sigma} \tilde{t}_{(13)\sigma}^{(2)\delta} f_{(13)}^{(a_2)} + \tilde{t}_{(a_2 1)}^{(2)\gamma\sigma} \tilde{t}_{(23)\sigma}^{(2)\delta} f_{(23)}^{(a_2)}] B_1(Q_{\psi\gamma X}), \quad (128)$$

$$\langle \gamma 1^{-+} | (f_2 \eta) 2 \rangle = q_\mu \epsilon_{\nu\beta\gamma\delta} K^\beta \tilde{t}_{(\eta f_2)}^{(2)\gamma\sigma} \tilde{t}_{(12)\sigma}^{(2)\delta} f_{(12)}^{(f_2)} B_1(Q_{\psi\gamma X}), \quad (129)$$

$$\langle \gamma 1^{-+} | (a_2 \pi) 2 \rangle = q_\mu \epsilon_{\nu\beta\gamma\delta} K^\beta [\tilde{t}_{(a_2 2)}^{(2)\gamma\sigma} \tilde{t}_{(13)\sigma}^{(2)\delta} f_{(13)}^{(a_2)} + \tilde{t}_{(a_2 1)}^{(2)\gamma\sigma} \tilde{t}_{(23)\sigma}^{(2)\delta} f_{(23)}^{(a_2)}] B_1(Q_{\psi\gamma X}). \quad (130)$$

For  $\psi \rightarrow \gamma 2^{++}$ , there are three independent couplings and two possible  $yz$  states,  $f_2 \eta$  and  $a_2 \pi$ .

$$\langle \gamma 2^{++} | (f_2 \eta) 1 \rangle = P_{\mu\nu\alpha\lambda}^{(2)}(K) \epsilon^{\alpha\beta\gamma\delta} K_\beta \tilde{t}_{(\eta f_2)}^{(1)} \tilde{t}_{(12)\delta}^{(2)\lambda} f_{(12)}^{(f_2)} \quad (131)$$

$$\langle \gamma 2^{++} | (a_2 \pi) 1 \rangle = P_{\mu\nu\alpha\lambda}^{(2)}(K) \epsilon^{\alpha\beta\gamma\delta} K_\beta [\tilde{t}_{(a_2 1)\gamma}^{(1)} \tilde{t}_{(23)\delta}^{(2)\lambda} f_{(23)}^{(a_2)} + \tilde{t}_{(a_2 2)\gamma}^{(1)} \tilde{t}_{(13)\delta}^{(2)\lambda} f_{(13)}^{(a_2)}], \quad (132)$$

$$\langle \gamma 2^{++} | (f_2 \eta) 2 \rangle = g_{\mu\nu} p_\psi^\lambda p_\psi^\sigma P_{\lambda\sigma\alpha\beta}^{(2)}(K) B_2(Q_{\psi\gamma X}) \epsilon_{\gamma\delta\alpha'}^\alpha K^\gamma \tilde{t}_{(\eta f_2)}^{(1)\delta} \tilde{t}_{(12)}^{(2)\alpha'\beta} f_{(12)}^{(f_2)}, \quad (133)$$

$$\begin{aligned} \langle \gamma 2^{++} | (a_2 \pi) 2 \rangle = & g_{\mu\nu} p_\psi^\lambda p_\psi^\sigma P_{\lambda\sigma\alpha\beta}^{(2)}(K) \epsilon_{\gamma\delta\alpha'}^\alpha K^\gamma [\tilde{t}_{(a_2 1)\delta}^{(1)} \tilde{t}_{(23)}^{(2)\alpha'\lambda} f_{(23)}^{(a_2)} \\ & + \tilde{t}_{(a_2 2)\delta}^{(1)} \tilde{t}_{(13)}^{(2)\alpha'\lambda} f_{(13)}^{(a_2)}] B_2(Q_{\psi\gamma X}), \end{aligned} \quad (134)$$

$$\langle \gamma 2^{++} | (f_2 \eta) 3 \rangle = q_\mu p_\psi^\lambda P_{\nu\lambda\alpha\beta}^{(2)}(K) B_2(Q_{\Psi\gamma X}) \epsilon_{\gamma\delta\alpha'}^\alpha K^\gamma \tilde{t}_{(\eta f_2)}^{(1)\delta} \tilde{t}_{(12)}^{(2)\alpha'\beta} f_{(12)}^{(f_2)}, \quad (135)$$

$$\langle \gamma 2^{++} | (a_2 \pi) 3 \rangle = q_\mu p_\psi^\lambda P_{\nu\lambda\alpha\beta}^{(2)}(K) \epsilon_{\gamma\delta\alpha'}^\alpha K^\gamma [\tilde{t}_{(a_2 1)\delta}^{(1)} \tilde{t}_{(23)}^{(2)\alpha'\beta} f_{(23)}^{(a_2)} + \tilde{t}_{(a_2 2)\delta}^{(1)} \tilde{t}_{(13)}^{(2)\alpha'\beta} f_{(13)}^{(a_2)}] B_2(Q_{\psi\gamma X}). \quad (136)$$

For  $\psi \rightarrow \gamma 2^{-+}$ , we have

$$\langle \gamma 2^{-+} | (f_0 \eta) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha \tilde{t}_{(f_0 \eta)}^{(2)\beta\gamma} q_\gamma f_{(12)}^{(f_0)} B_1(Q_{\psi\gamma X}), \quad (137)$$

$$\langle \gamma 2^{-+} | (a_0 \pi) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha \left\{ \tilde{t}_{(a_0 1)}^{(2)\beta\gamma} f_{(23)}^{(a_0)} + \tilde{t}_{(a_0 2)}^{(2)\beta\gamma} f_{(13)}^{(a_0)} \right\} q_\gamma B_1(Q_{\psi\gamma X}), \quad (138)$$

$$\langle \gamma 2^{-+} | (f_0 \eta) 2 \rangle = S_{\mu\nu} p_\psi^\gamma p_\psi^\delta \tilde{t}_{(f_0 \eta)}^{(2)\gamma\delta} f_{(12)}^{(f_0)} B_3(Q_{\psi\gamma X}), \quad (139)$$

$$\langle \gamma 2^{-+} | (a_0 \pi) 2 \rangle = S_{\mu\nu} p_\psi^\gamma p_\psi^\delta \left\{ \tilde{t}_{(a_0 1)}^{(2)\gamma\delta} f_{(23)}^{(a_0)} + \tilde{t}_{(a_0 2)}^{(2)\gamma\delta} f_{(13)}^{(a_0)} \right\} B_3(Q_{\psi\gamma X}), \quad (140)$$

$$\langle \gamma 2^{-+} | (f_0 \eta) 3 \rangle = q_\mu S_{\nu\gamma} \tilde{t}_{(f_0 \eta)}^{(2)\gamma\delta} p_\psi^\delta f_{(12)}^{(f_0)} B_3(Q_{\psi\gamma X}), \quad (141)$$

$$\langle \gamma 2^{-+} | (a_0 \pi) 3 \rangle = q_\mu S_{\nu\gamma} p_\psi^\delta \left\{ \tilde{t}_{(a_0 1)}^{(2)\gamma\delta} f_{(23)}^{(a_0)} + \tilde{t}_{(a_0 2)}^{(2)\gamma\delta} f_{(13)}^{(a_0)} \right\} B_3(Q_{\Psi\gamma X}), \quad (142)$$

$$\langle \gamma 2^{-+} | (f_2 \eta) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha q_\gamma P^{(2)\beta\gamma\beta'\gamma'}(K) \tilde{t}_{(12)\beta'\gamma'}^{(2)} f_{(12)}^{(f_2)} B_1(Q_{\psi\gamma X}) \quad (143)$$

$$\begin{aligned} \langle \gamma 2^{-+} | (a_2 \pi) 1 \rangle = & \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha P^{(2)\beta\gamma\beta'\gamma'}(K) q_\gamma B_1(Q_{\psi\gamma X}) [\tilde{t}_{(23)\beta'\gamma'}^{(2)} f_{(23)}^{(a_2)} + \tilde{t}_{(13)\beta'\gamma'}^{(2)} f_{(13)}^{(a_2)}], \\ & \end{aligned} \quad (144)$$

$$\langle \gamma 2^{-+} | (f_2 \eta) 2 \rangle = S_{\mu\nu} p_\psi^\gamma p_\psi^\delta P_{\gamma\delta\gamma'\delta'}^{(2)}(K) \tilde{t}_{(12)}^{(2)\gamma'\delta'} f_{(12)}^{(f_2)} B_3(Q_{\psi\gamma X}), \quad (145)$$

$$\langle \gamma 2^{-+} | (a_2 \pi) 2 \rangle = S_{\mu\nu} p_\psi^\gamma p_\psi^\delta P_{\gamma\delta\gamma'\delta'}^{(2)}(K) B_3(Q_{\psi\gamma X}) [\tilde{t}_{(23)}^{(2)\gamma'\delta'} f_{(23)}^{(a_2)} + \tilde{t}_{(13)}^{(2)\gamma'\delta'} f_{(13)}^{(a_2)}], \quad (146)$$

$$\langle \gamma 2^{-+} | (f_2 \eta) 3 \rangle = q_\mu S_{\nu\gamma} p_\psi^\delta P^{(2)\gamma\delta\gamma'\delta'}(K) \tilde{t}_{(12)\gamma'\delta'}^{(2)} f_{(12)}^{(f_2)} B_3(Q_{\Psi\gamma X}), \quad (147)$$

$$\langle \gamma 2^{-+} | (a_2 \pi) 3 \rangle = q_\mu S_{\nu\gamma} p_\psi^\delta P^{(2)\gamma\delta\gamma'\delta'}(K) B_3(Q_{\Psi\gamma X}) [\tilde{t}_{(23)\gamma'\delta'}^{(2)} f_{(23)}^{(a_2)} + \tilde{t}_{(13)\gamma'\delta'}^{(2)} f_{(13)}^{(a_2)}] \quad (148)$$

with  $S_{\mu\nu}$  defined as in Eq.(118).

### 3.3 $\psi \rightarrow \gamma K \bar{K} \pi$

Possible intermediate channels for this process are  $K^*K, K_0^*K, K_2^*K, a_0\pi, a_2\pi$ . The formulae for  $K_0^*K, K_2^*K, a_0\pi, a_2\pi$  intermediate states to the  $K\bar{K}\pi$  final state are the same as for the  $a_0\pi, a_2\pi, f_0\eta, f_2\eta$  intermediate states given in the previous subsection for the  $\pi^+\pi^-\eta$  final state. So here we only give partial wave amplitudes  $U_{\mu\nu}^i$  with  $K^*K$  intermediate states. We denote  $K, \bar{K}, \pi$  as particle 1,2,3.

$$\langle \gamma 0^{-+} | (K^*K) 1 \rangle = S_{\mu\nu} B_1(Q_{\psi\gamma X}) [\tilde{t}_{(K^*\bar{K})\lambda}^{(1)} \tilde{t}_{(13)}^{(1)\lambda} f_{(13)}^{(K^*)} + \tilde{t}_{(K^*\bar{K})\lambda}^{(2)} \tilde{t}_{(23)}^{(1)\lambda} f_{(23)}^{(K^*)}], \quad (149)$$

$$\langle \gamma 1^{++} | (K^*K) 1 \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha [\tilde{t}_{(23)}^{(1)\beta} f_{(23)}^{(K^*)} + \tilde{t}_{(13)}^{(1)\beta} f_{(13)}^{(K^*)}], \quad (150)$$

$$\langle \gamma 1^{++} | (K^*K) 2 \rangle = q_\mu S_{\nu\beta} B_2(Q_{\psi\gamma X}) [\tilde{t}_{(23)}^{(1)\beta} f_{(23)}^{(K^*)} + \tilde{t}_{(13)}^{(1)\beta} f_{(13)}^{(K^*)}], \quad (151)$$

$$\langle \gamma 1^{-+} | (K^*K) 1 \rangle = g_{\mu\nu} S_{\gamma\delta} [\tilde{t}_{(K^*\bar{K})}^{(1)\gamma} \tilde{t}_{(13)}^{(1)\delta} f_{(13)}^{(K^*)} + \tilde{t}_{(\bar{K}^*K)}^{(1)\gamma} \tilde{t}_{(23)}^{(1)\delta} f_{(23)}^{(K^*)}] B_1(Q_{\psi\gamma X}), \quad (152)$$

$$\langle \gamma 1^{-+} | (K^*K) 2 \rangle = q_\mu \epsilon_{\nu\beta\gamma\delta} K^\beta [\tilde{t}_{(K^*\bar{K})}^{(1)\gamma} \tilde{t}_{(13)}^{(1)\delta} f_{(13)}^{(K^*)} + \tilde{t}_{(\bar{K}^*K)}^{(1)\gamma} \tilde{t}_{(23)}^{(1)\delta} f_{(23)}^{(K^*)}] B_1(Q_{\psi\gamma X}). \quad (153)$$

### 3.4 $\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$

Listed here are formulae used in Refs.[8, 14].

$$\begin{aligned} \langle \gamma 0^{-+} | \rho\rho \rangle &= S_{\mu\nu} \epsilon_{\gamma\delta\lambda\sigma} p_1^\gamma p_2^\delta p_3^\lambda p_4^\sigma B_1(Q_{\psi\gamma X}) [f_{(12)}^{(\rho)} f_{(34)}^{(\rho)} B_1(Q_{\rho 12}) B_1(Q_{\rho 34}) \\ &\quad B_1(Q_{X(12)(34)}) - f_{(14)}^{(\rho)} f_{(32)}^{(\rho)} B_1(Q_{\rho 14}) B_1(Q_{\rho 32}) B_1(Q_{X(14)(32)})], \end{aligned} \quad (154)$$

$$\langle \gamma 0^{++} | \sigma\sigma \rangle = g_{\mu\nu} [f_{(12)}^{(\sigma)} f_{(34)}^{(\sigma)} + f_{(14)}^{(\sigma)} f_{(32)}^{(\sigma)}], \quad (155)$$

$$\begin{aligned} \langle \gamma 0^{++} | \rho\rho \rangle &= g_{\mu\nu} [f_{(12)}^{(\rho)} f_{(34)}^{(\rho)} (q_{(12)} \cdot q_{(34)}) B_1(Q_{\rho 12}) B_1(Q_{\rho 34}) + \\ &\quad f_{(14)}^{(\rho)} f_{(32)}^{(\rho)} (q_{(14)} \cdot q_{(32)}) B_1(Q_{\rho 14}) B_1(Q_{\rho 32})], \end{aligned} \quad (156)$$

$$\begin{aligned} \langle \gamma 0^{++} | \pi\pi'(\pi\sigma) \rangle &= g_{\mu\nu} [f_{(123)}^{(\pi')} (f_{(12)}^{(\sigma)} + f_{(32)}^{(\sigma)}) + f_{(234)}^{(\pi')} (f_{(23)}^{(\sigma)} + f_{(34)}^{(\sigma)}) + \\ &\quad f_{(143)}^{(\pi')} (f_{(14)}^{(\sigma)} + f_{(34)}^{(\sigma)}) + f_{(214)}^{(\pi')} (f_{(21)}^{(\sigma)} + f_{(14)}^{(\sigma)})], \end{aligned} \quad (157)$$

$$\begin{aligned} \langle \gamma 0^{++} | \pi\pi'(\pi\rho) \rangle &= g_{\mu\nu} [f_{(123)}^{(\pi')} f_{(12)}^{(\rho)} q_{(12)\alpha} (p_3 - p_{(12)})^\alpha B_1(Q_{\pi'\rho 3}) B_1(Q_{\rho 12}) + \\ &\quad f_{(234)}^{(\pi')} f_{(23)}^{(\rho)} q_{(23)\gamma} (p_4 - p_{(23)})^\gamma B_1(Q_{\pi'\rho 4}) B_1(Q_{\rho 23}) + \\ &\quad + \{1 \leftrightarrow 3\} + \{2 \leftrightarrow 4\} + \{1 \leftrightarrow 3 \ \& \ 2 \leftrightarrow 4\}], \end{aligned} \quad (158)$$

$$\begin{aligned} \langle \gamma 0^{++} | \pi a_1(\pi\rho) \rangle &= g_{\mu\nu} [P_{\alpha\beta}^{(1)} (p_{(123)}) p_4^\alpha q_{(12)}^\beta f_{(123)}^{(a_1)} f_{(12)}^{(\rho)} B_1(Q_{X a_1 4}) B_1(Q_{\rho 12}) \\ &\quad + P_{\alpha\beta}^{(1)} (p_{(234)}) p_1^\alpha q_{(23)}^\beta f_{(234)}^{(a_1)} f_{(23)}^{(\rho)} B_1(Q_{X a_1 1}) B_1(Q_{\rho 23})] \end{aligned}$$



$$+ \{1 \leftrightarrow 3\} + \{2 \leftrightarrow 4\} + \{1 \leftrightarrow 3 \& 2 \leftrightarrow 4\}], \quad (159)$$

$$\langle \gamma 2^{++} | (yy) 1 \rangle = X_{\mu\nu}^{(yy)}, \quad (160)$$

$$\langle \gamma 2^{++} | (yy) 2 \rangle = g_{\mu\nu} p_\psi^\alpha p_\psi^\beta X_{\alpha\beta}^{(yy)} B_2(Q_{\psi X\gamma}), \quad (161)$$

$$\langle \gamma 2^{++} | (yy) 3 \rangle = q_\mu X_{\nu\alpha}^{yy} p_\psi^\alpha B_2(Q_{\psi X\gamma}), \quad (162)$$

$$\langle \gamma 2^{++} | (f_2\sigma) 1 \rangle = P_{\mu\nu\alpha\beta}^{(2)}(K) \tilde{t}_{(12)}^{\alpha\beta} f_{(12)}^{(f_2)} f_{(34)}^{(\sigma)} + \{1 \leftrightarrow 3\} + \{2 \leftrightarrow 4\} + \{1 \leftrightarrow 3 \& 2 \leftrightarrow 4\}, \quad (163)$$

$$\langle \gamma 2^{++} | (f_2\sigma) 2 \rangle = g_{\mu\nu} p_\psi^\alpha p_\psi^\beta P_{\alpha\beta\gamma\delta}^{(2)}(K) \tilde{t}_{(12)}^{(2)\gamma\delta} f_{(12)}^{(f_2)} f_{(34)}^{(\sigma)} B_2(Q_{\psi X\gamma}) + \{1 \leftrightarrow 3\} \\ + \{2 \leftrightarrow 4\} + \{1 \leftrightarrow 3 \& 2 \leftrightarrow 4\}, \quad (164)$$

$$\langle \gamma 2^{++} | (f_2\sigma) 3 \rangle = q_\mu p_\psi^\beta P_{\nu\beta\gamma\delta}^{(2)}(K) \tilde{t}_{(12)}^{(2)\gamma\delta} f_{(12)}^{(f_2)} f_{(34)}^{(\sigma)} B_2(Q_{\psi X\gamma}) + \{1 \leftrightarrow 3\} \\ + \{2 \leftrightarrow 4\} + \{1 \leftrightarrow 3 \& 2 \leftrightarrow 4\}. \quad (165)$$

The amplitudes involving X particles of  $J^P = 2^{++}$  involves a rank two tensor,  $X^{(yy)}$ . The definition of this is given below:

$$X_{\mu\nu}^{(\sigma\sigma)} = f_{(12)}^{(\sigma)} f_{(34)}^{(\sigma)} B_2(Q_{X(12)(34)}) P_{\mu\nu\alpha\beta}^{(2)}(K) (p_{(12)}^\alpha - p_{(34)}^\alpha) (p_{(12)}^\beta - p_{(34)}^\beta) + \{2 \leftrightarrow 4\} \quad (166)$$

$$X_{\mu\nu}^{(\rho\rho)} = f_{(12)}^{(\rho)} f_{(34)}^{(\rho)} B_1(Q_{\rho 12}) B_1(Q_{\rho 34}) P_{\mu\nu\alpha\beta}^{(2)}(K) \tilde{t}_{(12)}^{(1)\alpha} \tilde{t}_{(34)}^{(1)\beta} + \{2 \leftrightarrow 4\} \quad (167)$$

where  $L = 2$  decay for  $X \rightarrow \rho\rho$  is ignored in view of the centrifugal barrier suppression.

From the flux tube model for hybrids,  $1^{-+}$  hybrids with  $I = 0$  decay dominantly into  $4\pi$  through  $a_1\pi$ . Then the  $\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$  is an ideal place for finding  $1^{-+}$  hybrids. With high statistics data at BES and CLEO-C, one should look for the iso-scalar  $1^{-+}$  hybrid in this channel. Here we add the formulae for  $1^{-+}$  hybrid production.

$$\langle \gamma 1^{-+} | [\pi a_1(\rho\pi)] 1 \rangle = g_{\mu\nu} p_\psi^\alpha P_{\alpha\beta}^{(1)}(K) [P^{(1)\beta\gamma}(p_{(123)}) \tilde{t}_{(12)\gamma}^{(1)} f_{(123)}^{(a_1)} f_{(12)}^{(\rho)} \\ + P^{(1)\beta\gamma}(p_{(234)}) \tilde{t}_{(23)\gamma}^{(1)} f_{(234)}^{(a_1)} f_{(23)}^{(\rho)}] + \{1 \leftrightarrow 3\} + \{2 \leftrightarrow 4\} \\ + \{1 \leftrightarrow 3 \& 2 \leftrightarrow 4\}, \quad (168)$$

$$\langle \gamma 1^{-+} | [\pi a_1(\rho\pi)] 2 \rangle = q_\mu P_{\nu\beta}^{(1)}(K) [P^{(1)\beta\gamma}(p_{(123)}) \tilde{t}_{(12)\gamma}^{(1)} f_{(123)}^{(a_1)} f_{(12)}^{(\rho)} \\ + P^{(1)\beta\gamma}(p_{(234)}) \tilde{t}_{(23)\gamma}^{(1)} f_{(234)}^{(a_1)} f_{(23)}^{(\rho)}] + \{1 \leftrightarrow 3\} + \{2 \leftrightarrow 4\} \\ + \{1 \leftrightarrow 3 \& 2 \leftrightarrow 4\}. \quad (169)$$

### 3.5 $\psi \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

We construct the amplitudes  $U_{\mu\nu}^i$  with a notation similar to the previous subsection for the  $\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$  channel. Here we denote  $K^+, K^-, \pi^+, \pi^-$  as 1,2,3,4.

$$\langle \gamma 0^{-+} | K^* \bar{K}^* \rangle = S_{\mu\nu} \epsilon_{\gamma\delta\lambda\sigma} p_1^\gamma p_2^\delta p_3^\lambda p_4^\sigma f_{(14)}^{(K^*)} f_{(23)}^{(\bar{K}^*)}$$

$$B_1(Q_{\psi\gamma X})B_1(Q_{K^*14})B_1(Q_{\bar{K}^*23})B_1(Q_{X(14)(23)}), \quad (170)$$

$$\langle \gamma 0^{++} | \kappa \kappa \rangle = g_{\mu\nu} f_{(14)}^{(\kappa)} f_{(23)}^{(\kappa)}, \quad (171)$$

$$\langle \gamma 0^{++} | K^* \bar{K}^* \rangle = g_{\mu\nu} \tilde{t}_{(14)}^{(1)\alpha} \tilde{t}_{(23)\alpha}^{(1)} f_{(14)}^{(K^*)} f_{(23)}^{(K^*)}, \quad (172)$$

$$\langle \gamma 0^{++} | K^* \kappa \rangle = g_{\mu\nu} [\tilde{t}_{(K^*\bar{\kappa})\alpha}^{(1)} \tilde{t}_{(14)}^{(1)\alpha} f_{(14)}^{(K^*)} f_{(23)}^{(\kappa)} + \tilde{t}_{(\bar{K}^*\kappa)\alpha}^{(1)} \tilde{t}_{(23)}^{(1)\alpha} f_{(23)}^{(K^*)} f_{(14)}^{(\kappa)}], \quad (173)$$

$$\langle \gamma 1^{++} | (K^* \bar{K}^*) 1 \rangle = \epsilon_{\mu\nu\lambda\alpha} p_\psi^\lambda \epsilon^{\alpha\beta\gamma\delta} K_\beta \tilde{t}_{(14)\gamma}^{(1)} \tilde{t}_{(23)\delta}^{(1)} f_{(14)}^{(K^*)} f_{(23)}^{(\bar{K}^*)}, \quad (174)$$

$$\langle \gamma 1^{++} | (K^* \bar{K}^*) 2 \rangle = q_\mu S_{\nu\alpha} \epsilon^{\alpha\beta\gamma\delta} K_\beta \tilde{t}_{(14)\gamma}^{(1)} \tilde{t}_{(23)\delta}^{(1)} f_{(14)}^{(K^*)} f_{(23)}^{(\bar{K}^*)} B_2(Q_{\psi\gamma X}), \quad (175)$$

$$\begin{aligned} \langle \gamma 1^{++} | (K^* \kappa) 1 \rangle &= \epsilon_{\mu\nu\lambda\alpha} p_\psi^\lambda \epsilon^{\alpha\beta\gamma\delta} p_{X\beta} [\tilde{t}_{(K^*\bar{\kappa})\gamma}^{(1)} \tilde{t}_{(14)\delta}^{(1)} f_{(14)}^{(K^*)} f_{(23)}^{(\kappa)} \\ &\quad + \tilde{t}_{(\bar{K}^*\kappa)\gamma}^{(1)} \tilde{t}_{(23)\delta}^{(1)} f_{(23)}^{(K^*)} f_{(14)}^{(\kappa)}], \end{aligned} \quad (176)$$

$$\begin{aligned} \langle \gamma 1^{++} | (K^* \kappa) 2 \rangle &= q_\mu S_{\nu\alpha} \epsilon^{\alpha\beta\gamma\delta} p_{X\beta} [\tilde{t}_{(K^*\bar{\kappa})\gamma}^{(1)} \tilde{t}_{(14)\delta}^{(1)} f_{(14)}^{(K^*)} f_{(23)}^{(\kappa)} \\ &\quad + \tilde{t}_{(\bar{K}^*\kappa)\gamma}^{(1)} \tilde{t}_{(23)\delta}^{(1)} f_{(23)}^{(K^*)} f_{(14)}^{(\kappa)}] B_2(Q_{\psi\gamma X}), \end{aligned} \quad (177)$$

$$\langle \gamma 2^{++} | (yz) 1 \rangle = X_{\mu\nu}^{(yz)}, \quad (178)$$

$$\langle \gamma 2^{++} | (yz) 2 \rangle = g_{\mu\nu} p_\psi^\alpha p_\psi^\beta X_{\alpha\beta}^{(yz)} B_2(Q_{\psi\gamma X}), \quad (179)$$

$$\langle \gamma 2^{++} | (yz) 3 \rangle = q_\mu p_\psi^\alpha X_{\alpha\nu}^{(yz)} B_2(Q_{\psi\gamma X}), \quad (180)$$

$$\langle \gamma 4^{++} | (yy) 1 \rangle = Z_{\mu\nu\lambda\sigma}^{(yy)} p_\psi^\lambda p_\psi^\sigma B_2(Q_{\psi\gamma X}), \quad (181)$$

$$\langle \gamma 4^{++} | (yy) 2 \rangle = g_{\mu\nu} p_\psi^\alpha p_\psi^\beta p_\psi^\gamma p_\psi^\delta Z_{\alpha\beta\gamma\delta}^{(yy)} B_4(Q_{\psi\gamma X}), \quad (182)$$

$$\langle \gamma 4^{++} | (yy) 3 \rangle = q_\mu Z_{\nu\lambda\sigma\alpha}^{(yy)} p_\psi^\lambda p_\psi^\sigma p_\psi^\alpha B_4(Q_{\psi\gamma X}), \quad (183)$$

$$\langle \gamma 2^{-+} | (K^* \bar{K}^*) 1A \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha A^{\beta\lambda} p_{\psi\lambda} B_1(Q_{\psi\gamma X}), \quad (184)$$

$$\langle \gamma 2^{-+} | (K^* \bar{K}^*) 1B \rangle = \epsilon_{\mu\nu\alpha\beta} p_\psi^\alpha B^{\beta\lambda} p_{\psi\lambda} B_1(Q_{\psi\gamma X}), \quad (185)$$

$$\langle \gamma 2^{-+} | (K^* \bar{K}^*) 2A \rangle = S_{\mu\nu} A^{\lambda\sigma} p_{\psi\lambda} p_{\psi\sigma} B_3(Q_{\psi\gamma X}), \quad (186)$$

$$\langle \gamma 2^{-+} | (K^* \bar{K}^*) 2B \rangle = S_{\mu\nu} B^{\lambda\sigma} p_{\psi\lambda} p_{\psi\sigma} B_3(Q_{\psi\gamma X}), \quad (187)$$

$$\langle \gamma 2^{-+} | (K^* \bar{K}^*) 3A \rangle = q_\mu S_{\nu\gamma} A^{\gamma\delta} p_{\psi\delta} B_3(Q_{\psi\gamma X}), \quad (188)$$

$$\langle \gamma 2^{-+} | (K^* \bar{K}^*) 3B \rangle = q_\mu \epsilon_{\nu\gamma} B^{\gamma\delta} p_{\psi\delta} B_3(Q_{\psi\gamma X}) \quad (189)$$

with  $S_{\mu\nu}$  defined as Eq.(118). The amplitudes involving X particles of  $J^P = 2^+$  involves a rank two tensor,  $X^{(yz)}$ . The definition of this is given below:

$$X_{\mu\nu}^{(\kappa\kappa)} = \tilde{t}_{(\kappa\bar{\kappa})\mu\nu}^{(2)} f_{(14)}^{(\kappa)} f_{(23)}^{(\kappa)}, \quad (190)$$

$$X_{\mu\nu}^{(K^* \bar{K}^*)} = P_{\mu\nu\alpha\beta}^{(2)}(K) \tilde{t}_{(14)}^{(1)\alpha} \tilde{t}_{(23)}^{(1)\beta} f_{(14)}^{(K^*)} f_{(23)}^{(\bar{K}^*)}, \quad (191)$$

$$X_{\mu\nu}^{(K^* \kappa)} = P_{\mu\nu\alpha\beta}^{(2)}(K) [\tilde{t}_{(K^*\bar{\kappa})\alpha}^{(1)} \tilde{t}_{(14)}^{(1)\beta} f_{(14)}^{(K^*)} f_{(23)}^{(\kappa)} + \tilde{t}_{(\bar{K}^*\kappa)\alpha}^{(1)} \tilde{t}_{(23)}^{(1)\beta} f_{(23)}^{(K^*)} f_{(14)}^{(\kappa)}], \quad (192)$$

$$Z_{\alpha\beta\gamma\delta}^{(\kappa\kappa)} = \tilde{t}_{(\kappa\bar{\kappa})\alpha\beta\gamma\delta}^{(4)} f_{(14)}^{(\kappa)} f_{(23)}^{(\kappa)}, \quad (193)$$

$$Z_{\alpha\beta\gamma\delta}^{(K^*\bar{K}^*)} = P_{\alpha\beta\gamma\delta\alpha'\beta'\gamma'\delta'}^{(4)}(K)\tilde{t}_{(K^*\bar{K}^*)}^{(2)\alpha'\beta'}\tilde{t}_{(14)}^{(1)\gamma'}\tilde{t}_{(23)}^{(1)\delta'}f_{(14)}^{(K^*)}f_{(23)}^{(\bar{K}^*)}, \quad (194)$$

$$A_{\mu\nu} = P_{\mu\nu\alpha\alpha'}^{(2)}(K)\epsilon^{\alpha\beta\gamma\delta}K_{\beta}\tilde{t}_{(14)\gamma}^{(1)}\tilde{t}_{(23)\delta}^{(1)}\tilde{t}_{(K^*\bar{K}^*)}^{(1)\alpha'}f_{(14)}^{(K^*)}f_{(23)}^{(\bar{K}^*)}, \quad (195)$$

$$B_{\mu\nu} = P_{\mu\nu\alpha\alpha'}^{(2)}(K)\epsilon^{\alpha\beta\gamma\delta}K_{\beta}\tilde{t}_{(K^*\bar{K}^*)\gamma}^{(1)}(\tilde{t}_{(14)\delta}^{(1)}\tilde{t}_{(23)}^{(1)\alpha'} + \tilde{t}_{(23)\delta}^{(1)}\tilde{t}_{(14)}^{(1)\alpha'})f_{(14)}^{(K^*)}f_{(23)}^{(\bar{K}^*)} \quad (196)$$

where  $A_{\mu\nu}$  corresponds to  $2^{-+} \rightarrow K^*\bar{K}^*$  with  $L = 1$  and  $S = 1$ ,  $B_{\mu\nu}$  corresponds to  $2^{-+} \rightarrow K^*\bar{K}^*$  with  $L = 1$  and  $S = 2$ . We ignore  $2^{-+} \rightarrow K^*\bar{K}^*$  with  $L = 3$  due to a strong centrifugal barrier.

## 4 Discussion

Here we add some points of general technique in fitting data. The first concerns the fact that tensor amplitudes are not always unique. As an example, in  $J/\psi \rightarrow \gamma f_2$ , there are three independent helicity amplitudes. But the general formalism allows one to write down five covariant tensor amplitudes. Those five are independent in the process  $J/\psi \rightarrow \omega f_2$ , but for the radiative decay, gauge invariance makes two of them dependent on the other three. Two further linear combinations differ from the first three only by different  $s$ -dependence arising from the momentum dependence built into the tensor expressions. Chung [11] recommends using all five combinations, so as to retain the differences in possible  $s$ -dependence. However, this gives rise to a practical problem.

One is usually fitting resonances such as the  $f_2$  to data. If two of the amplitudes differ from the others only in  $s$ -dependence, this is equivalent to putting into the numerator of an  $f_2$  Breit-Wigner amplitude a linear combination of two  $s$ -dependent terms with two free parameters. This may lead to a zero amplitude at the resonance mass and can give rise to structure which may lie 500 MeV or 1 GeV away from the  $f_2$ ; it may be easily confused with the effects of other resonances. This is illustrated in Fig.1 for the amplitude squared  $|T|^2$  taking as an example  $J/\psi \rightarrow \gamma f_2(1700) \rightarrow \gamma K\bar{K}$ . For the solid line, we use  $T = Q_{\psi\gamma f}^2 B_2(Q_{\psi\gamma f})/(M_f^2 - s - iM_f\Gamma_f)$  with  $M_f = 1.7\text{GeV}$  and  $\Gamma_f = 0.15\text{GeV}$ ; for the dotted line which lies very close to the solid line, we use  $T = 30.3 * Q_{\psi\gamma f}^4 B_4(Q_{\psi\gamma f})/(M_f^2 - s - iM_f\Gamma_f)$ . The two different  $s$ -dependence numerators give a hardly visible difference in line shape of  $f_2$ . But if one allows two  $s$ -dependent terms in the numerator with two free parameters, the ridiculous shape (dashed line) could happen for a single resonance  $f_2(1700)$ ; in this illustration we use has  $20[Q_{\psi\gamma f}^2 B_2(Q_{\psi\gamma f}) - 30.3Q_{\psi\gamma f}^4 B_4(Q_{\psi\gamma f})]$  in the numerator. Although theoretically this possibility cannot be excluded, it is very odd and in practice one may end up fitting other  $2^{++}$  components far away from the  $f_2$  resonance mass with the  $f_2$ . One therefore should be very careful in drawing conclusions from a fit using more than the minimum number of amplitudes with different angular dependence.

In  $J/\psi$  radiative decays, the  $c\bar{c}$  pair annihilates to gluons. This requires a short-range interaction with a range of order  $1/m_c$ , where  $m_c$  is the mass of the  $c$  quark. Therefore




Figure 1: Distortion on Breit-Wigner amplitude squared by the s-dependence numerators: with  $Q_{\psi\gamma f}^2 B_2(Q_{\psi\gamma f})$  (solid line), with  $30.3Q_{\psi\gamma f}^4 B_4(Q_{\psi\gamma f})$  (dotted line),  $20[Q_{\psi\gamma f}^2 B_2(Q_{\psi\gamma f}) - 30.3Q_{\psi\gamma f}^4 B_4(Q_{\psi\gamma f})]$  (dashed line).

the centrifugal barrier for  $J/\psi \rightarrow \gamma X$  is strong. Some production with  $L = 1$  is observed (at momentum transfer  $\leq 1$  GeV/c), but we find little evidence for  $L > 1$ .

## 5 Acknowledgement

We thank the Royal Society for funds allowing a collaboration between Queen Mary College, University of London and IHEP, Beijing. We thank V.V.Anisovich and A.V.Sarantsev for many helpful discussions. A small part of formulae presented here were independently written and cross checked by them. We thank L.Y.Dong, Z.J.Guo and J.L.Hu for Monte Carlo simulation of some formulae in this paper and for pointing out some typos. The work is partly supported by CAS Knowledge Innovation Project (KJ CX2-SW-N02) and the National Natural Science Foundation of China under Grant No.10225525.

## References

- [1] R.A.Briere et al., Preprint CLNS-01-1742.
- [2] V.V.Anisovich, D.V.Bugg, A.V.Sarantsev, Nucl. Phys. A537 (1992) 501; V.V.Anisovich et al., Nucl. Phys. A544 (1992) 747.
- [3] A.V.Anisovich et al., Phys. Lett. B452 (1999) 173; Phys. Lett. B452 (1999) 180; Phys. Lett. B452 (1999) 187; Phys. Lett. B468 (1999) 304; Phys. Lett. B468 (1999) 309; Phys. Lett. B500 (2001) 222; Phys. Lett. B507 (2001) 23.
- [4] A.V.Anisovich, V.A.Sadovnikova, Euro. Phys. J. A2 (1998) 199; A.V.Anisovich et al., J. Phys. G28 (2002) 15.
- [5] BES Collaboration, Phys. Lett. B440 (1998) 217.
- [6] BES Collaboration, Phys. Lett. B446 (1999) 356.
- [7] BES Collaboration, Phys. Lett. B472 (2000) 200.
- [8] BES Collaboration, Phys. Lett. B472 (2000) 208.
- [9] BES Collaboration, Phys. Lett. B476 (2000) 25.
- [10] W.Rarita, J.Schwinger, Phys. Rev. **60**, 61 (1941); E.Behtends and C.Fronsdal, Phys. Rev. **106**, 345 (1957).
- [11] S.U.Chung, Phys. Rev. **D48**, 1225 (1993); Phys. Rev. **D57**, 431 (1998).

- [12] F. von Hippel and C.Quigg, Phys. Rev. **D5**, 624 (1972).
- [13] W.Greiner et al., Quantum Chromodynamics, Springer 2002, p.242.
- [14] D.V.Bugg et al., Phys. Lett. **B353**, 378 (1995).